

## Genetic Relationships among *Aegilops* L. and *Triticum* L. Species based on the Internal Transcribed Spacer Sequences of nrDNA (ITS)

<sup>1</sup>Lina Mammdouh Alnaddaf, <sup>2</sup>Mohammad Yahia Moualla and <sup>3</sup>Nadia Haider

<sup>1,2</sup>Department of Field Crops, Faculty of Agriculture, Field Crops Tishreen University, Lattakia, Syria

<sup>3</sup>Department of Molecular Biology and Biotechnology, Atomic Energy Commission, Damascus, Syria

**Abstract:** Studying of genetic relationships among *Triticum* L. and *Aegilops* L. species by direct sequencing of PCR-amplified Internal Transcribed Spacer (ITS) of nuclear ribosomal DNA to investigate the polymorphism in nucleotide sequences among 19 *Aegilops* L. and 7 *Triticum* L. species. ITS sequences were aligned with CLUSTAL W 2.1 multiple sequence alignment program. The phylogenetic relationships among species were reconstructed using Unweighted Pair Group Mean Arithmetic average (UPGMA) and Neighbor-Joining (NJ) methods. ITS region ranged from 600 to 602 bp. The length of ITS1 was 221-222 bp and ITS2 was 215-217 bp. The 5.8S subunit was 163 bp long. The G+C content of the ITS1 region ranged from (60.3 to 63.9) % in all *Triticum* and *Aegilops* species. The G+C content of the 5.8S subunit was entirely identical in all species, 59.5%. The G+C content of the ITS2 region ranged from 59.9 to 63.5%. There were 67 variable sites (11.12%) in the entire ITS region. The phylogenetic relationships among species were reconstructed using Unweighted Pair Group Mean Arithmetic average (UPGMA) were discussed.

**Keywords:** *Aegilops*, genetic relationships, internal transcribed spacer, ITS, *Triticum*

### INTRODUCTION

Ribosomal RNA genes (known as ribosomal DNA or (rDNA) are found as parts of repeat units that are arranged in tandem arrays, located at the chromosomal sites known as Nucleolar Organizing Regions (NORs). Each repeat unit consists of a transcribed region (having genes for 18S, 5.8S and 26S rRNAs and the external transcribed spacers i.e., ETS1 and ETS2) and a Non-Transcribed Spacer (NTS) region. In the transcribed region, Internal Transcribed Spacers (ITS) are found on either side of 5.8S rRNA gene and are described as ITS1 and ITS2 (Sharma *et al.*, 2002). These regions can be readily amplified by PCR and sequenced using universal primers (White *et al.*, 1990). Universal PCR primers designed from highly conserved regions flanking the ITS and its relatively small size (600-700 bp) enable easy amplification of ITS region due to high copy number (up to-30000 per cell) (Dubouzet and Shinoda, 1999) of rDNA repeats. However, DNA from many plant species may not be amplified by the currently available primers and thus species-specific ITS primers would be desirable (Lin *et al.*, 2010). there is now a set of seven ITS universal primers (White *et al.*, 1990; Sun *et al.*, 1994) that are widely applied for diagnostic (Linder *et al.*, 2000) and phylogenetic studies in many angiosperm families (Baldwin *et al.*,

1995; Haider, 2011). Because of their different rates of evolution, ITS regions presented high variability and faster rates of evolution (Baldwin *et al.*, 1995; Barker *et al.*, 1988; Kollipara *et al.*, 1997; Downie and Kartz-Downie, 1996) and could be highly applicable in a broad range of organisms including plants, fungi and animals (White *et al.*, 1990). The ITS polymorphism might occur at the genus, species or individual levels, making it to be a useful source of molecular characters for phylogenetic studies in many angiosperms (Nwakanma *et al.*, 2003). This makes the ITS region an interesting subject for evolutionary/phylogenetic investigations (Baldwin *et al.*, 1995; Hershkovitz and Zimmer, 1996; Hershkovitz *et al.*, 1999; Nwakanma *et al.*, 2003; Gulbitti-Onarici *et al.*, 2009) as well as biogeographic investigations (Baldwin, 1993; Suh *et al.*, 1993; Hsiao *et al.*, 1994; Dubouzet and Shinoda, 1999; Sharma *et al.*, 2002). ITS loci has properties that were claimed to be advantageous for purposes of phylogenetic reconstruction: Biparental inheritance-Universality-Simplicity -Intragenomic uniformity-Intergenomic variability-Low functional (González-Chavira *et al.*, 2006). In the last decade numerous molecular markers and techniques were used for studies on origin, evolution and relationships in the wheat group: chloroplast and nuclear microsatellite markers (Lelley *et al.*, 2000; Ishii *et al.*, 2001), chromosome-

**Corresponding Author:** Lina Mammdouh Alnaddaf, Department of Field Crops, Faculty of Agriculture, Field Crops Tishreen University, Lattakia, Syria

specific low-copy DNA (Liu *et al.*, 2003), nuclear genes (Caldwell *et al.*, 2004) and ITS has also been used to study the evolution of wheat species in the early 1980s (Peacock *et al.*, 1981; Dvorák and Appels, 1982; Wang *et al.*, 2000; Rudnóy *et al.*, 2005). The evolution of the ITS region is more complicated in hybrid and polyploid species (Baldwin *et al.*, 1995; Wendel *et al.*, 1995; Waters and Schaal, 1996). Baldwin *et al.* (1995) proposed that ITS sequences would provide direct evidence of reticulate evolution if concerted evolution failed to homogenize the repeat units contributed by different parental species when the hybridization event was recent, or if nrDNA repeats were at different loci in the parental genomes and interlocus gene conversion was inoperative in their hybrid, or if the hybrid was asexual. Since the history of polyploid wheats is relatively short (Mori *et al.*, 1995) and the ITS repeats in polyploid wheats are located at different loci (Dubkovsky and Dvorák, 1995; Badaeva *et al.*, 1996), it may be possible to identify ITS sequences of different parental origins in polyploid wheats and hence to identify their progenitors. In addition, the *Triticum* complex is a good model system for studying how hybridization and polyploidization could possibly affect the evolution of nrDNA (Zhang *et al.*, 2002). Numerous studies have demonstrated the utility of the ITS region (Hsiao *et al.*, 1995a; Wang *et al.*, 2000; Blattner, 2004; Jakob *et al.*, 2010) for resolving relationships among closely related species in *Triticeae* and other plant species (Hsiao *et al.*, 1994, 1995a, b; Baldwin *et al.*, 1995; Goel *et al.*, 2002; Sharma *et al.*, 2002; Zhang *et al.*, 2002; Alvarez and Wendel, 2003; Bordbar *et al.*, 2011). Hsiao *et al.* (1995a) studied the sequence of the ITS region of 30 diploid *Triticeae* species representing 19 genomes. They suggest that the sequence of the ITS is variable enough to differentiate closely related species (Hsiao *et al.*, 1994). Nalini *et al.* (2007) found out that the polymorphic ITS fragments were correlated with Quantitative Trait Loci (QTL) such as the spike size and the number of spikelets per spike. And the polymorphic ITS fragments detected there should be further cloned and sequenced in order to screen for interesting agronomic traits with potential use in wheat breeding. Carvalho *et al.* (2009a) In their previous study performed in 48 old Portuguese bread wheat cultivars, they found out nine ITS PCR-RFLP patterns with the enzymes *AluI*, *HpaII* and *TaqI* (three patterns per enzyme). ITS PCR-RFLP markers revealed a lower percentage of polymorphism when compared with other DNA markers such as ISSRs (Carvalho *et al.*, 2009b). Carvalho *et al.* (2011) In their present study, 51 durum wheat cultivars showed 40% of ITS variation and this lower level of polymorphism could be due to the absence of the D genome. Low ITS variation was previously reported for *Triticeae* (Zhang *et al.*, 2002)

and other taxa such as *Cucurbitaceae* (Jobst *et al.*, 1998); *Oleaceae* (Jeandroz *et al.*, 1997) and *Vigna* (Saini *et al.*, 2008). So Carvalho *et al.* (2011) concluded that the knowledge of the genetic relationships and phylogenies among the durum wheat cultivars and their botanical varieties might contribute for the designing of intraspecific crosses between the genotypes studied there, with potential interest for wheat improvement. The sequence data of the ITS region has also been studied earlier to assess genetic diversity in cultivated barley (Petersen and Seberg, 1996). Sharma *et al.* (2002) used Internal Transcribed Spacer (ITS) sequences of ribosomal DNA of wild barley and their comparison with ITS sequences in common wheat to assess genetic diversity at the intraspecific level in barley and wheat.

Variations in length and sequence of the ITS have been used for taxonomic study in trees (Campbell *et al.*, 1995; Baldwin, 1992; Lin *et al.*, 2010) used ITS region sequence to identify phylogenetic analysis in intraspecific population of the genus *Zanthoxylum*. Sun *et al.* (1994) studied the phylogenetic relationships of the genus *Sorghum* and related genera by sequencing the ITS region. Their results were consistent with those based on chloroplast DNA analysis (Cao, 1997). Moore and Field (2005) were able to identify the roots of different plant species in mixed samples using sequences of the nuclear ITS region. Sequencing of the same region proved useful for DNA barcoding of members of the *Cycadales*. Singh *et al.* (2008) were used Internal Transcribed Spacer (ITS) sequences of nuclear ribosomal DNA to assess genetic diversity and phylogenetic relationships in chickpea (*C. arietinum*) cultivars and its related wild species. Their present comprehensive study has provided novel insights in the phylogeny of genus *Cicer*. Linder *et al.* (2000) developed a method that uses (ITS) region of the nuclear ribosomal DNA (rDNA) repeat to identify most underground plant parts to the level of species. Their technique should allow the below-ground parts of plants in any system to be identified and thereby open new possibilities for the study of below-ground plant communities.


In this study we study the genetic relationships among *Triticum* and *Aegilops* species inferred from the sequence data of the ITS region of the nuclear rDNA to broaden the genetic base of wheat and improve its yield, quality and resistance to biotic and abiotic stresses.

## MATERIALS AND METHODS

**Plant materials:** The plant material consisted of 26 accessions (Table 1). 19 accessions representing (19) *Aegilops* L species: *Ae. crassa*, *Ae. tauschii*, *Ae. umbellulata*, *Ae. triuncialis*, *Ae. comosa*, *Ae. peregrine*, *Ae. caudata*, *Ae. biuncialis*, *Ae. ovata*, *Ae. neglecta*, *Ae. speltoides*, *Ae. ventricosa*, *Ae. searsii*, *Ae. cylindrica*, *Ae. kotschyi*, *Ae. longissima*, *Ae. bicornis*, *Ae.*

Table 1: Base compositions of ITS1, 5.8S and ITS2 regions of *Triticum* and *Aegilops* species

<i>Triticum</i> and <i>Aegilops</i> species	ITS1	% CG	5.8	% CG	ITS2	% CG	ITS total	% CG	
<i>Ae. bicornis</i>	25_R_primer-5R	222	61.7	163	59.5	217	63.1	602	61.4
<i>Ae. sharonensis</i>	26_R_primer-5R	222	61.7	163	59.5	217	63.1	602	61.4
<i>Ae. crassa</i>	27_R_primer-5R	222	61.2	163	59.5	216	62.6	601	61.4
<i>Ae. triuncialis</i>	28_R_primer-5R	222	60.7	163	59.5	217	63.1	602	61.4
<i>Ae. peregrina</i>	29_R_primer-5R	222	60.3	163	59.5	217	62.2	602	60.6
<i>Ae. biuncialis</i>	30_R_primer-5R	222	63.0	163	59.5	217	62.2	602	61.5
<i>Ae. ovata</i>	31_R_primer-5R	222	63.0	163	59.5	217	62.2	602	61.5
<i>Ae. neglecta</i>	32_R_primer-5R	222	63.5	163	59.5	217	62.6	602	61.8
<i>Ae. searsii</i>	33_R_primer-5R	222	61.5	163	59.5	217	63.1	602	61.3
<i>Ae. longissima</i>	34_R_primer-5R	222	62.5	163	59.5	217	63.1	602	61.7
<i>Ae. tauschii</i>	35_R_primer-5R	222	61.2	163	59.5	217	60.3	602	60.3
<i>Ae. umbellulata</i>	36_R_primer-5R	222	61.2	163	59.5	217	63.1	602	61.2
<i>Ae. comosa</i>	37_R_primer-5R	222	62.6	163	59.5	217	61.7	602	61.2
<i>Ae. caudata</i>	38_R_primer-5R	222	60.7	163	59.5	217	63.1	602	61.1
<i>Ae. uniaristata</i>	39_R_Rev-Primer	222	61.5	163	59.5	217	63.1	602	61.3
<i>Ae. speltoides</i>	40_R_Rev-Primer	222	61.5	163	59.5	217	63.5	602	61.5
<i>T. turgidum</i>	41_R_Rev-Primer	221	61.5	163	59.5	217	60.8	601	60.6
<i>T. dicoccon</i>	42_R_Rev-Primer	222	61.2	163	59.5	217	63.1	602	61.2
<i>T. aestivum</i>	43_R_Rev-Primer	222	61.2	163	59.5	217	60.8	602	60.6
<i>Ae. ventricosa</i>	44_R_Rev-Primer	222	62.6	163	59.5	217	59.9	602	60.6
<i>Ae. cylindrica</i>	45_R_Rev-Primer	222	63.9	163	59.5	217	60.3	602	61.2
<i>Ae. kotschyi</i>	46_R_Rev-Primer	222	61.2	163	59.5	217	63.1	602	61.2
<i>T. monococcum</i>	48_R_Rev-Primer	222	61.2	163	59.5	215	61.3	600	60.6
<i>T. urartu</i>	25_Rev_5R-Rev	222	61.2	163	59.5	217	61.2	602	60.6
<i>T. dicoccoides</i>	26_Rev_5R-Rev	222	61.5	163	59.5	217	60.6	602	60.5
<i>T. durum</i>	27_Rev_5R-Rev	222	61.5	163	59.5	217	60.6	602	60.5

Fig. 1: PCR of ITS in *Aegilops* species

M: 100 bp DNA; 25: *Ae. bicornis*; 26: *Ae. sharonensis*; 27: *Ae. crassa*; 28: *Ae. triuncialis*; 29: *Ae. peregrina*; 30: *Ae. biuncialis*; 31: *Ae. ovata*; 32: *Ae. neglecta*

*uniaristata*, *Ae. sharonensis*. (7) accessions representing 7 *Triticum* L. species: *T. monococcum*, *T. urartu*, *T. dicoccoides*, *T. durum*, *T. turgidum*, *T. dicoccon*, *T. aestivum* (Table 1). All the accessions were obtained from the Genetic Resources Unit (GRU) at the International Centre for Agricultural Research in the Dry Areas (ICARDA), Aleppo, Syria.

**DNA extraction:** Total genomic DNA was isolated from fresh, young leaves as described by Doyle and Doyle (1987). DNA was quantified using Spectrometer and the concentration of all samples was set at 10 ng/ $\mu$ L.

**PCR amplification:** PCR reactions were carried out in a 25  $\mu$ L volume containing 10 $\times$ PCR buffer (Eurobio), dNTPs (10 mM) (Mix Roche), 10 $\times$ MgC 12 (50 mM) (Eurobio), Taq polymerase (5 U/ $\mu$ L) (Eurobio). ITS region (including ITS 1, 5.8 S and ITS 2) was amplified using the following primer pair (White *et al.*, 1990):

ITS-4 (5'-TCCTCCGCTTATTGATATGC-3') -ITS-5 (5'GGAAGTAAAAGTCGTAACAAGG-3'), DNA was added to each PCR at a rate of 10 ng and the total volume was adjusted with dd H<sub>2</sub>O to 25  $\mu$ L. For 35 cycles, PCRs were subjected to 95°C for 1min for DNA denaturation, 50°C for 1 min for annealing of primers, 72°C for 1 min for extension of the target region and 72°C for 5 min for final extension. PCR products (1-5  $\mu$ L) were digested according to manufacturer (Fermentaz). Digested fragments were separated by electrophoresis on 1.8% agarose gel that was run at 100 V for 2 h in TBE 0.5 x buffer and visualised under UV lights (Fig. 1).

**Sequence analyses:** The PCR products delivered to Leibniz University Hannover-Department of Plant Biotechnology-Hannover-Germany for sequencing, which was conducted in both forward and reverse directions using the amplification primers. ITS sequences were aligned with CLUSTAL W 2.1 multiple sequence alignment program. The boundaries of the ITS region was determined by comparison with the sequence information in (Chatterton *et al.*, 1992). The phylogenetic relationships among species were reconstructed using Unweighted Pair Group Mean Arithmetic average (UPGMA) and Neighbor-Joining (NJ) methods.

## RESULTS AND DISCUSSION

Earlier studies in the *Poaceae* have focused on morphology, anatomy, taxonomy, physiology, cytology, genetics and crop improvement. They have provided important information, but data based on these

studies are not enough to assess the true relationships between these species. Phylogenetic constructions proposed for the *Triticum* and *Aegilops* species based on these characters are poorly resolved and differ widely in topology. Therefore we used (ITS) regions of the nuclear ribosomal DNA (nrDNA) because it has been shown to be a valuable source of evidence to resolve phylogenetic relationships in many angiosperm groups (Gulbitti-Onarici *et al.*, 2009).

The results of this study showed that all *Triticum* L. and *Aegilops* L. species presented a 700-bp PCR product of invariant length (Fig. 1). And The ITS region ranged from 600 to 602 bp. The length of ITS1 was 221-222 bp and ITS2 was 215-217 bp. The 5.8S subunit was 163 bp long (Table 1). The total length of the entire ITS of rDNA among *Triticum* L. and *Aegilops* L. species during the present study were variable and were in agreement with the results of earlier studies. Hsiao *et al.* (1995a) studied the phylogenetic relationships of 30 diploid species of *Triticeae* representing 19 genomes based on the analysis of the ITS region of the 18-26S rRNA gene. The length of the entire ITS region of the *Triticeae* species studied only varied from 596 to 605 nucleotides. In diploid species of the *Triticeae*, the ITS1 region ranges from 216 to 223 bp in length and the ITS2 ranges from 214 to 217 base pairs. Our results were similar to the research of Cao (1997) who used the internal transcribed spacer to determine the phylogenetic relationship of hexaploid wheats among and within the five groups of hexaploid wheat. The ITS 1, 5.8S and ITS2 regions in the five groups of hexaploid wheat contained 222, 163 and 217 bp, respectively. Both spacer regions (ITS1, ITS2) were identical in length to those of Chinese Spring (Chatterton *et al.*, 1992), but there was a 1% sequence difference in the spacer regions between Chinese Spring (Chatterton *et al.*, 1992) and the common wheat cultivar Columbus (Cao, 1997). Also Wang *et al.* (2000) studied nine diploid species representing six sections of *Aegilops* using direct sequencing of PCR-amplified DNA fragments for ITS region. The ITS region ranged from 601 to 607 bp. The length of ITS1 was 221-226 bp and ITS2 was 215-217 bp. The 5.8S subunit sequence was the most conserved region and was 164 bp long in all the diploid species sequenced. Furthermore, no nucleotide substitution was observed in this subunit. This is agree with our results. In addition Zhang *et al.* (2002) Studied the origin and evolution of tetraploid wheats based on the Internal Transcribed Spacer (ITS) sequences of nuclear ribosomal DNA. The results of their study showed that 2-bp indel was found in ITS1 sequences and 1-bp indel in ITS2 sequences. and The length ranged from 221 to 223 bp in ITS1 and from 216 to 217 bp in ITS2. This is agree in part with our results, there is a 1-bp indel in ITS1 sequences of *T. turgidum* at position 211 and an 3-bp indel in ITS2 sequences of *Ae. crassa* at position (416) and *T. monococcum* at position (510-511). Also Bordbar *et al.* (2011) observed

The difference occurred in the length of ITS2, which ranged from 214 to 216 bp. That difference was due to one dinucleotide (AA/TT) deletion at positions 185-186 of ITS2 in some sequences of *Ae. crassa*, *Ae. vavilovii* and *Ae. juvenalis*.

Based on the results of this study The G+C content of the ITS1 region ranged from (60.3 to 63.9) % in all *Triticum* and *Aegilops* species. The G+C content of the 5.8S subunit was entirely identical in all species, 59.5%. The G+C content of the ITS2 region ranged from 59.9 to 63.5% (Table 1). But this G+C content were differ according to Wang *et al.* (2000) mentioned that The GC content of the *Aegilops* species was fairly equivalent, 61.1-62.9%. so The G+C contents of the sequences in the study of Zhang *et al.* (2002) were varied from 59 to 62% for ITS1 and from 60 to 65% for ITS2. In addition, Nalini *et al.* (2007) reported in identification and characterization of some ITS variants from hexaploid wheat (*Triticum aestivum* L.) The length of the three ITS variants were found to be same but showed variation in their %G+C. Generally %G+C was slightly more in ITS1 (range: 59.00-60.36) than in ITS2 (range: 60.82-61.75). on the other hand Asay *et al.* (1992) observed that mountain rye and wheat (*Triticum aestivum*) have complete homology in the 5.8 s gene but have different nucleotides at 13 and 30 locations in the ITS1 and ITS2 regions, respectively. nevertheless, the AT 39% and GC 61% contents are the same as those in wheat. that is in contrast with 25% AT and 75% GC in rice. The ITS region (601 bp) in mountain rye is 1bp shorter than that in bread wheat (602) and 4 bp longer than that in barley (*Hordeum vulgare*).

The available studies have indicated that may be there is a relationship between the comparatively high GC content and adaptation to the environment. however it is evident that if ITSs are subject to select pressure, temperature is only one factor (Torres *et al.*, 1990). This hypothesis certainly is preliminary and needs to be further tested.

Our present study showed that There were 67 variable sites (11.12%) in the entire ITS region. The ITS1 region had 28 variable sites and the ITS2 region had 39 variable sites. In the 5.8 s gene there were no variable sites. And that 16 of 62 variable sites were in *T. dicoccon*, *Ae. kotschyi*, *T. durum* and one in *Ae. biuncialis*. whereas, Sharma *et al.* (2002) indicated that The sequence data showed that the total variable sites in wild barley (7.4%) were comparable with the total variable sites in wheat (6.90%). Also Cao (1997) reported that there were 13 variable sites (2%) in the ITS1 and ITS2 regions among the five groups of hexaploid wheat.

In the current study, among the substitutions, transitions were more frequent than transversions, the transitions in ITS1 were 17 while in ITS2 were 22. the transversions were 10 in both ITS1, ITS2 and among the indels, deletions were one in ITS1 and 7 in ITS2 (Table 2).

Table 2: The gray bases indicate transition, transversion, deletion

30_R_primer-5R	TCTATTTAAT
27_Rev_5R-Rev	GCTATTTAAT
41-R_Rev-Primer	-CTATAAAT

```

TCGTGACCCT GACCAAAACA GACCGCGCAC GCGTCATCCA ATCCGTCGGC
GACGGCATCG TCCGTGCTC GGCCAATGCC TCGACCACCT CCCCTCCTGG
GAGCGGGTGG GGGCTCGGGG TAAAAGAACC CACGGCGCCA AAGGCGTCAA
GGAAACACTGT GCCTAACCCG GGGGCATGGC TAGCTTGCTA GCCGTCCCTC
GTGTTGAAA GCTATTTAAT CCACAGACT CTCGGCAACG GATATCTCGG
CTCTCGCATC GATGAAGAAC GTAGCGAAAT GCGATACCTG GTGTGAATTG
CAGAATCCCG CGAACCATCG AGTCTTTGAA CGCAAGTTGC GCCCGAGGCC
ACTCGGCCGA GGGCAGCCT GCCTGGGCGT CACGCCAAAA CACGCTCCCA
ACCACCCTCA TCGGGAATCG GGATGCGGCA TCTGGTCCCT CGTCTCGCAA
GGGGCGGTGG ACCGAAGATC GGGCTGCCGG TGTACCGCGC CGGACACAGC
GCATGGTGGG CFTCTCGCT TTATCAACGC AGTGATCCG ACGCGACGC
GGCATTATGG CCTCAGAATG ACCCAGCAA CGAAGCGCAT GTCGCTTCGA
CC

```

Fig. 2: The ITS sequence of *Ae. bicornis* the highly conserved area within the ITS1 is indicated in bold

However, the frequency of such sites in the present study was higher than those reported earlier in the ITS sequences of different species of *Triticeae*. For instance, in a study involving ten genotypes of cultivated barley, *H. vulgare*, only 0.84% substitution sites were available while the indels were completely absent (Petersen and Seberg, 1996). In another study involving different species of *Triticeae* including wild and cultivated barley, no substitutions or indels were detected in a study of two accessions of each species procured from widely separated geographical regions (Hsiao *et al.*, 1995a).

A characteristic conserved sequence GGCRY- (4 to 7n) GYGCAAGGAA (where Y = C or T, R = G or A), was also available in the ITS1 of both *Triticum* L. and *Aegilops* L. species (Fig. 2). In previous studies on many flowering plants this characteristic sequence has been reported in the middle of ITS1 and this sequence is presumed as a recognition site for processing of a primary transcript into the structural rRNA (Liu and Schardl, 1994). So we can notice this conserved sequence in the middle of ITS1 at position (134-150) (GGCGCCGAAGGCGTCAAGGAA) and we observe 2 transversions (G→A) at position 140 in two *Aegilops* species *Ae. bicornis*, *Ae. searsii*.

The phylogenetic tree of 26 *Triticum* and *Aegilops* species generated using the Unweighted Pair Group Mean Arithmetic Average (UPGMA) and Neighbor-Joining (NJ) methods was shown in (Fig. 3). two trees are essentially identical. There were mainly three clades in this tree. The first clade included *T. turgidum*, *T. aestivum*, *T. dicoccon*, *T. durum*. The second clade consisted of *Ae. speltoides*, *Ae. triuncialis*, *Ae. caudata*, *Ae. ovata*, *Ae. biuncialis*, *Ae. umbellulata*, *Ae. peregrina*, *Ae. bicornis*, *Ae. sharonensis*, *Ae. comosa*, *Ae. neglecta*, *Ae. tauschii*, *Ae. cylindrica*, *Ae. ventricosa*. The third clade contained of *T. dicoccoides*, *T. monococcum*, *Ae. uniaristata*, *Ae. searsii*, *Ae. longissima*, *Ae. crassa*, *Ae. kotschy*, *T. urartu* (Fig. 3).

The wheats (*Triticum* spp.) form a polyploid series with diploid (2n = 2x = 14), tetraploid (2n = 4x = 28) and hexaploid (2n = 6x = 42) forms. The diploid wheats comprise a single genomic group with the genome formula AA (*T. monococcum*, *T. urartu*). The tetraploid emmer wheats are divided into two groups, those with the genome formula AABB (*T. turgidum*) and those with the genome formula AAGG (*T. timopheevi*). On evidence it appears that the wild allotetraploid emmer wheat *T. turgidum* ssp. *dicoccoides* (AABB) arose by amphyploidy between the wild diploid wheat *T. urartu* (AA) and a diploid member of the *Aegilops* genus (BB) (Rudnóy *et al.*, 2002). The origin of B genome is still a matter of debate. Polyphyletic origin or divergent evolution of B genome from the donor species are hypothesized. On the basis of chondriome divergence *Ae. speltoides* seems to be the cytoplasm donor (female parent) of the tetraploid wheats (Wang *et al.*, 2000). A descendant of *T. turgidum* ssp. *dicocoides*, the *T. turgidum* ssp. *dicoccon* was probably the ancient tetraploid from which hexaploid wheats (AABBDD) may have evolved by hybridisation between the AABB tetraploid as cytoplasm donor and the D genome diploid *Ae. taushii*. (Huang *et al.*, 2002). Our results in the first clade reflect these facts (Fig. 3). the current study showed that *Ae. speltoides* was separated from remaining four *Sitopsis* species.

Zhang *et al.* (2002) reported that Phylogenetic analysis demonstrated that *Aegilops speltoides* was distinct from other species in *Aegilops* sect. *Sitopsis*. Also Sliai and Amer (2011) revealed that *Ae. speltoides* does not form a monophyletic clade with other *Sitopsis* species (Goriunova *et al.*, 2008; Salina *et al.*, 2006). Recent studies showed that *Ae. speltoides* was the main contributor of the B genome of polyploid wheats (Huang *et al.*, 2002). In addition the sequence of one chloroplast gene (*rbcL*, for the Rubisco large subunit) from seven *Triticum* and *Aegilops* species indicated that *Ae. speltoides* is the donor of both the plasmon and B genome of common wheat (Terachi *et al.*, 1988; Wang *et al.*, 1997; Gupta *et al.*, 2008; Al-ahmar *et al.*, 2010). And the tree reconstructed based on data of ten EST-SSRs mapped on the B genome showed that *Ae. speltoides* had the closest relationship with *T. aestivum* and *T. durum* (Zhang *et al.*, 2006). Yen *et al.* (2005) observed the cytoplasm of *T. turgidum* L. is very similar to that in some races of *Ae. speltoides*.

In our ITS analysis, a close relationship of the sequences from *Ae. tauschii* (*Ae. squarrosa*) and *Ae. cylindrica* was found. Wang *et al.* (1997) reported that *Ae. squarrosa* is the maternal parent of three tetraploids, *Ae. cylindrica*, *Ae. crassa* and *Ae. ventricosa* (Wan *et al.*, 2002). Also according to Queen *et al.* (2004) The three D genome-containing *Aegilops* species, comprising the two members of Section *Vertebrata* (*Ae. ventricosa*, *Ae. tauschii*; Van Slageren




Fig. 3: Phenogram of UPGMA cluster analysis in *Triticum* L. and *Aegilops* L. species

1994) and *Ae. cylindrica* form a clade. More over Kharazian (2008) said *Ae. taushii* and *Ae. cylindrica* have similar genome which it grouped together in the application of Rf data (the migration distance of the band/distance of solvent front) (Jaaska, 1981, 1993) but in the MW (the molecular weight of prolamin bands) are separated. Wang *et al.* (2000) provide an approach to understand the genome evolution of allopolyploid species in genus *Aegilops* through studied the Evolution of parental ITS regions of nuclear rDNA in allopolyploid *Aegilops* (*Poaceae*) species and assumed that they might be largely homogenized by concerted evolution toward one of their other ancestors during the process of hybridization and polyploidization. These evidences have been observed in this study. Based in this conclusion we observed a close relationships between (*Ae. tauschii* and *Ae. cylindrica*), (*Ae. triuncialis* and *Ae. caudata*), (*Ae. comosa* and *Ae. neglecta*), (*Ae. biuncialis*, *Ae. ovata*, *Ae. peregrina* and *Ae. umbellulata*). Wang *et al.* (1997) indicated that *Ae. triuncialis* arose from *Ae. umbellulata* as mother and their results supports the dimaternal origin of *Ae. triuncialis* from reciprocal crosses between *Ae. umbellulata* and *Ae. caudata*. Konstantinos and Bebeli (2010) observed *Ae. triuncialis* (genome UC) grouped in the same subgroup with *Ae. caudata* (C), which is its progenitor male parent. on the other hand they noticed that, *Ae. peregrina*-SU grouped closer to the male parent (*Ae. umbellulata*-U) than the female parent (species from *Sitopsis* section possibly *Ae. searsii*-S). In addition our ITS sequence-based phylogeny showed *Ae. ovata*, *Ae. triuncialis*, *Ae. peregrina*, *Ae. biuncialis*

formed a clade with their common ancestor, *Ae. umbellulata* (UU) (Fig. 3). Wang *et al.* (2000) pointed out that *Ae. umbellulata* (UU) and *Ae. ovata* (UUMM) have the closest relationship among the polyploids with U genome. their results were in agreement with the similarities of ecological and morphological characters of these two species. Wang *et al.* (1997) reported that *Ae. biuncialis* arose from *Ae. umbellulata* as mother.

In the second clade there was two sister group (*Ae. comosa*, *Ae. neglecta*) and (*Ae. sharonensis*, *Ae. bicornis*) (Fig. 3). Mendlinger and Zohary (1995) reported that *Ae. sharonensis* was found to be equally close to *Ae. bicornis*. Wang *et al.* (1997) likewise observed the plasmon of *Ae. umbellulata* is closely related to the plasmons of *Ae. comosa*. and *Ae. neglecta* origin from reciprocal crosses between *Ae. umbellulata* and *Ae. comosa*.

*T. urartu*, *T. monococcum* belong to the *Einkorn* wheat group (Mizumoto *et al.*, 2002). These two species proposed as the A genome doner to polyploid wheats (Dvorák *et al.*, 1993; Takumi *et al.*, 1993). Our results revealed that there were Two different types of ITS sequences were found in *Triticum* tetraploid species. One type (*T. dicoccoides*) formed a group with *T. monococcum*. and the other type (*T. turgidum*) consisted of a group with other *Triticum* species. This finding is supported by Zhang *et al.* (2002) that mentioned the A genome of *T. dicoccoides* originated from *T. monococcum* based on the ITS 2 sequences of nuclear ribosomal DNA. The similarity between *T. dicoccoides* and *T. monococcum* could be the result of a recent introgression event (Fig. 3).

Tsunewaki and Ogihara (1983) noted that S<sup>b</sup> and S' plasma types found in *sitopsis* section showed much closer relation to A plasma type of *Einkorn* wheat than to other plasma types (B, G and S) of the same section. More over according to Wang *et al.* (1997) based on two trees (illustrates the phylogenetic trees constructed by Unweighted Pair-Group Method Using Arithmetic Averages (UPGMA) and Neighbor-Joining (NJ) methods) showed that *Einkorn* is closer to *Aegilops* than to *Triticum*. A similar result was reported by Cencki *et al.* (2008) using Random Amplification of Polymorphic DNA (RAPD) analysis to estimate the phylogenetic relationships among wild species of *Triticum* and *Aegilops* and cultivars of *Triticum aestivum* and *Triticum turgidum*. And this is agree with present study *T. urartu*, *T. monococcum* had a closer relationships with *Ae. uniaristata*, *Ae. searsii*, *Ae. longissima*, *Ae. crassa*, *Ae. kotschyi*. Terachi *et al.* (1984) observed a very close resemblance between *Ae. crassa* and *T. monococcum*. Whereas, Wang *et al.* (1997) mentioned that *Ae. searsii* was mostly similar to *Ae. kotschyi* and *Ae. kotschyi* arose from *Ae. searsii* as mother. The present results support this conclusion.

### CONCLUSION

- ITS 2 was more variable than ITS 1.
- *T. dicoccon*, *Ae. kotschyi*, *T. durum* were more variable than other species.
- *Einkorn* is closer to *Aegilops* than to *Triticum*.
- *Ae. ovata*, *Ae. triuncialis*, *Ae. peregrina*, *Ae. biuncialis* formed a clade with their common ancestor, *Ae. umbellulata* (UU).
- The similarity between *T. dicoccoides* and *T. monococcum* could be the result of a recent introgression event.
- Conserved area was identical in the ITS 1 of both *Triticum* and *Aegilops* species except two *Aegilops* species *Ae. bicornis*, *Ae. searsii*.
- Transitions were more frequent than transversions and the deletions were one in ITS 1 and 3 in ITS 2.
- The frequency of variable sites in the present study was higher than those reported earlier in the ITS sequences of different species of *Triticeae*.
- We can identify ITS sequences of different parental origins in *Triticum* and *Aegilops* species and hence to identify their progenitors.
- ITS used for resolving relationships among closely related species in *Triticeae* and other plant species.

### ACKNOWLEDGMENT

We would like to thank Dr. Nizar MirAli, Head of Department, for his support. We also thank I. Nabulsi, Q. Ghanemah, M. A. Kahla, E. Fawaz, N. Charfalden, M. C. Alden Laboratory of Molecular Biology and Biotechnology-Department of Molecular Biology and Biotechnology, Atomic Energy Commission, Damascus, Syria.

### REFERENCES

- Al-ahmar, E., N. Haider and H. Azzam, 2010. Genetic Relationships among *Aegilops* L. species using DNA molecular markers. Ph.D. Thesis, General Commission for Scientific Agricultural Research, Gene Bank Division, Faculty of Agriculture, Damascus University, Syria.
- Alvarez, I. and J.F. Wendel, 2003. Ribosomal ITS sequences and plant phylogenetic inference. *Mol. Phylogenet. Evol.*, 29: 435-455.
- Asay, K.H., C. Hsiao, N.J. Chatterton, K.B. Jensen and R.R.C. Wang, 1992. Nucleotide sequence of the internal transcribed spacer region of rDNA in mountain rye, *secale montanum* GUS. (Gramineae). *Plant Molecul. Biol.*, 20: 161-162.
- Badaeva, E., B. Friebe and B. Gill, 1996. Genome differentiation in *Aegilops*. 2. Physical mapping of 5S and 18S-26S ribosomal RNA gene families in diploid species. *Genome*, 39: 1150-1158.
- Baldwin, B.G., 1992. Phylogenetic utility of the internal transcribed spacers of nuclear ribosomal DNA in plants: An example from the compositae. *Mol. Phylogenet. Evol.*, 1: 3-16.
- Baldwin, B.G., 1993. Molecular phylogenetics of *Calycadenia* (Compositae) based on ITS sequences of nuclear ribosomal DNA: Chromosomal and morphological evolution reexamined. *Am. J. Bot.*, 80: 222-238.
- Baldwin, B.G., M.J. Sanderson, J.M. Porter, M.F. Wojciechowski, C.S. Campbell and M.J. Donoghue, 1995. The ITS region of nuclear ribosomal DNA: A valuable source of evidence on angiosperm phylogeny. *Ann. Mol. Bot. Gard.*, 82: 247-277.
- Barker, R.F., N.P. Harberd, M.G. Jarvist and R.B. Flavell, 1988. Structure and evolution of the intergenic region in a ribosomal DNA repeat unit of wheat. *J. Mol. Biol.*, 201: 1-17.
- Blattner, F.R., 2004. Phylogeny of *Hordeum* (Poaceae) as inferred by nuclear rDNA ITS sequences. *Mol. Phylogenet. Evol.*, 33: 289-299.
- Bordbar, F., M.R. Rahiminejad, H. Saeidi and R.F. Blattner, 2011. Phylogeny and genetic diversity of D-genome species of *Aegilops* and *Triticum* (Triticeae, Poaceae) from Iran based on microsatellites, ITS and trnL-F. *Plant Syst. Evol.*, 291: 117-131.
- Caldwell, K., J. Dvorak, E.S. Lagudah, E. Akhunov, M.C. Luo, P. Wolters and W. Powell, 2004. Sequence polymorphism in polyploid wheat and their d-genome diploid ancestor. *Genetics*, 167: 941-947.
- Campbell, C.S., M.F. Wojciechowski, B.G. Baldwin, L.A. Alice and M.J. Donoghue, 1995. Persistent nuclear ribosomal DNA sequence polymorphism in the *Amelanchier* agamic complex (Rosaceae). *Mol. Bio. Evol.*, 14: 81-90.

- Cao, W., 1997. Wheat taxonomy and cultivar identification using molecular markers. A Thesis Submitted to the Faculty of Graduate Studies and Research. Department of Crop Science and Plant Ecology, University of Saskatchewan.
- Carvalho, A., H. Guedes-Pinto and J. Lima-Brito, 2009a. Genetic variability assessed by ITS PCR-RFLP markers in old Portuguese bread wheat. *J. Genet.*, 88: 363-367.
- Carvalho, A., J. Lima-brito, B. Maças and H. Guedes-Pinto, 2009b. Genetic diversity and variation among botanical varieties of old Portuguese wheat cultivars revealed by ISSR assays. *Biochem. Genet.*, 47: 276-294.
- Carvalho, A., H. Guedes-Pinto and J. Lima-Brito, 2011. Physical localization of NORs and ITS length variants in old Portuguese durum wheat cultivars. *Indian Acad. Sci. J. Genet.*, 90(1).
- Cenkci, S., M. Yildiz, M. Konuk and Y. Eren, 2008. RAPD analyses of some wild *Triticum L.* and *Aegilops L.* species and wheat cultivars in Turkey. *Acta Biol. Cracovien. Botan.*, 50(1): 35-42.
- Chatterton, N.J., C. Hsiao, K.H. Asay, R.R.C. Wang and K.B. Jensen, 1992. Nucleotide sequence of the internal transcribed spacer region of rDNA in wheat, *Triticum aestivum L.* (Gramineae). *Plant Mol. Biol.*, 20: 59-160.
- Downie, S.R. and D.S. Kartz-Downie, 1996. A molecular phylogeny of Apiaceae subfamily Apioideae: evidence from nuclear ribosomal DNA internal transcribed spacer sequences. *Am. J. Bot.*, 83: 234-251.
- Doyle, J.J. and J.L. Doyle, 1987. A rapid DNA isolation procedure for small quantities of fresh leaf. *Phytochem. Bull.*, 19: 11-15.
- Dubkovsky, J. and J. Dvorák, 1995. Ribosomal RNA multigene loci: Nomads of the Triticeae genomes. *Genetics*, 140: 1367-1377.
- Dubouzet, J.G. and K. Shinoda, 1999. Relationships among old and New world Alliums according to ITS DNA sequence analysis. *Theor. Appl. Genet.*, 98: 422-433.
- Dvorák, J. and R. Appels, 1982. Chromosomal and nucleotide sequence differentiation in genomes of polyploid *Triticum* species. *Theor. Appl. Genet.*, 63: 349-360.
- Dvorák, J., P.D. Terlizzi, H.B. Zhang and P. Resta, 1993. The evolution of polyploid wheats: Identification of the A genome donor species. *Genome*, 36: 21-31.
- Goel, S., S.N. Raina and Y. Ogihara, 2002. Molecular evolution and phylogenetic implications of internal transcribed spacer sequences of nuclear ribosomal DNA in the Phaseolus-Vigna complex. *Mol. Phylogenet. Evol.*, 22: 1-19.
- González-Chavira, M.M., I.P. Torres, E.V. Pineda and R.G. Guevara-Gonzalez, 2006. DNA markers. *Adv. Agric. Food Biotechnol.*, 2006: 99-134.
- Goriunova, S.V., N.N. Chikida and E.Z. Kochieva, 2008. Molecular analysis of the phylogenetic relationships among the diploid *Aegilops* species of the section sitopsis. *Genetika*, 44(1): 137-141.
- Gulbitti-Onarici, S., C. Sancak, S. Sumer and S. Ozcan, 2009. Phylogenetic relationships of some wild wheat species based on the internal transcribed spacer sequences of nrDNA. *Curr. Sci.*, 96: 794-800.
- Gupta, P.K., R.R. Mir, A. Mohan and J. Kumar, 2008. Wheat genomics: Present status and future prospects. *Int. J. Plant Genomics.*, Article 896451, Doi: 10.1155/2008/896451.
- Haider, N., 2011. Chloroplast-specific universal primers and their uses in plant studies. *Biol. Plantarum.*, 55(2): 225-236
- Hershkovitz, M.A. and E.A. Zimmer, 1996. Conservation patterns in angiosperm rDNA ITS2 sequences. *Nucl. Acid Res.*, 24: 2857-2867.
- Hershkovitz, M.A., E.A. Zimmer and W.J. Hahn, 1999. Ribosomal DNA Sequences and Angiosperm Systematics. Taylor and Francis, London, pp: 268-326.
- Hsiao, C., N.J. Chatterton, K.H. Asay and K.B. Jensen, 1994. Phylogenetic relationships of 10 grass species: An assessment of phylogenetic utility of the internal transcribed spacer region in nuclear ribosomal DNA in monocots. *Genome*, 37: 112-120.
- Hsiao, C., N. Chatterton, K. Asay and K. Jensen, 1995a. Phylogenetic relationships of the monogenomic species of the wheat tribe Triticeae (Poaceae) inferred from nuclear rDNA (ITS) sequences. *Genome*, 38: 211-223.
- Hsiao, C., N. Chatterton, K. Asay and K. Jensen, 1995b. Molecular phylogeny of the Pooideae (Poaceae) based on nuclear rDNA (ITS) sequences. *Theor. Appl. Genet.*, 90: 389-398.
- Huang, S., A. Sirikhachornkit, X. Su, J. Faris, B. Gill, R. Haselkorn and P. Gornicki, 2002. Genes encoding plastid acetyl-CoA carboxylase and 3-phosphoglycerate kinase of the *Triticum/Aegilops* complex and the evolutionary history of polyploid wheat. *Proc. Natl. Acad. Sci. USA*, 99: 8133-8138.
- Ishii, T., N. Mori and Y. Ogihara, 2001. Evaluation of allelic diversity at chloroplast microsatellite loci among common wheat and its ancestral species. *Theor. Appl. Genet.*, 103: 896-904.
- Jaaska, V., 1981. Aspartate aminotransferase and alcohol dehydrogenase isoenzymes: Intraspecific differentiation in *Aegilops tauschii* and origin of the D genome polyploids in the wheat group. *Plant Syst. Evol.*, 137(4): 259-273.

- Jaaska, V., 1993. Isoenzymes in the Evaluation of Germplasm Diversity in Wild Diploid Relatives of Cultivated Wheat. In: Damania, A.B. (Ed.), Biodiversity and Wheat Improvement. John Wiley and Sons, New York, pp: 274- 257.
- Jakob, S.S., C. Heibl, D. Rödder and F.R. Blattner, 2010. Population demography influences climatic niche evolution: Evidence from diploid South American *Hordeum* species (Poaceae). *Mol. Ecol.*, 19: 1423-1438.
- Jeandroz, S., A. Roy and J. Bousquet, 1997. Phylogeny and phylogeography of the circumpolar genus *Fraxinus* (Oleaceae) based on internal transcribed spacer sequences of nuclear ribosomal DNA. *Mol. Phylogenet. Evol.*, 7: 241-251.
- Jobst, J., K. King and V. Hemleben, 1998. Molecular evolution of the internal transcribed spacers (ITS1 and ITS2) and phylogenetic relationships among species of the family Cucurbitaceae. *Mol. Phylogenet. Evol.*, 9: 204-219.
- Kharazian, N., 2008. Chemotaxonomic studies on *Aegilops* L. (Poaceae) in Iran. *Pak. J. Biol. Sci.*, 11(9): 1204-1211.
- Kollipara, K.P., R.J. Singh and T. Hymowitz, 1997. Phylogenetic and genomic relationships in the genus *Glycine* Willd. based on sequences from the ITS region of nuclear rDNA. *Genome*, 40: 57-68.
- Konstantinos, G.T. and P.J. Bebeli, 2010. Genetic diversity of Greek *Aegilops* species using different types of nuclear genome markers. *Molecul. Phylogenet. Evol.*, 56: 951-961.
- Lelley, T., M. Stachel, H. Grausgruber and J. Vollmann, 2000. Analysis of relationships between *Aegilops tauschii* and the D genome of wheat utilizing microsatellites. *Genome*, 43: 661-668.
- Lin, S.Y., W.P. Geun, K.O. Woung and H.S. Kwan, 2010. The internal transcribed spacer rDNA specific markers for identification of *Zanthoxylum piperitum*. *Afr. J. Biotechnol.*, 9(37): 6027-6039.
- Linder, C.R., L.A. Moore and R.B. Jackson, 2000. A universal molecular method for identifying underground plant parts to species. *Molecul. Ecol.*, 1549-1559.
- Liu, J.S. and C.L. Schardl, 1994. A conserved sequence in internal transcribed spacer 1 of plant nuclear rRNA genes. *Plant Mol. Biol.*, 26: 775-778.
- Liu, B., G. Segal, J.K. Rong and M. Feldman, 2003. A chromosome-specific sequence common to the B genome of polyploid wheat and *Aegilops searsii*. *Plant Syst. Evol.*, 241: 55-66.
- Mendlinger, S. and D. Zohary, 1995. The extent and structure of genetic variation in species of the *Sitopsis* group of *Aegilops*. *Heredity*, 74: 616-627.
- Mizumoto, K., S. Hirosawa, C. Nakamura and S. Takumi, 2002. Nuclear and chloroplast genome genetic diversity in the wild einkorn wheat, *Triticum urartu*, revealed by AFLP and SSLP analyses. *Hereditas*, 137: 208-214.
- Moore, L.A. and C.B. Field, 2005. A technique for identifying the roots of different species in mixed samples using nuclear ribosomal DNA. *J. Veg. Sci.*, 16: 131-134.
- Mori, N., Y. Liu and K. Tsunewaki, 1995. Wheat phylogeny determined by RFLP analysis of nuclear DNA wild tetraploid wheats. *Theor. Appl. Genet.*, 90: 129-134.
- Nalini, E., S.G. Bhagwat and N. Jawali, 2007. Identification and characterization of some ITS variants from hexaploid wheat (*Triticum aestivum* L.). *Plant Sci.*, 173: 262-268.
- Nwakanma, D.C., M. Pillay, B.E. Okoli and A. Tenkouano, 2003. PCR-RFLP of the ribosomal DNA Internal Transcribed Spacers (ITS) provides markers for the A and B genomes in *Musa* L. *Theor. Appl. Genet.*, 108: 154-159.
- Peacock, W.J., W.L. Gerlach and E.S. Dennis, 1981. Molecular Aspects of Wheat Evolution: Repeated DNA Sequences. Cambridge University, Cambridge, pp: 41-60.
- Petersen, G. and O. Seberg, 1996. ITS regions highly conserved in cultivated barleys. *Euphytica*, 90: 233-234.
- Queen, R.A., B.M. Gribbon, C. James, P. Jack and A.J. Flavell, 2004. Retrotransposon-based molecular markers for linkage and genetic diversity analysis in wheat. *Mol. Gen. Genomics.*, 271: 91-97.
- Rudnóy, S., Z. Bratek, E. Paldi, I. Racz and D. Lasztity, 2002. Chloroplast 16S rRNA sequences from different *Triticum* species. *Proc. 7th Hungarian Congress Plant Physiol.*, 46(3-4): 47-48.
- Rudnóy, S., Z. Bratek, E. Páldi, I. Rácz and D. Lásztity, 2005. Studies on chloroplast and nuclear rDNA in hexaploid bread wheat and its relatives. *Proc. 8th Hungarian Congress Plant Physiol. 6th Hungarian Conf. Photosynthesis*, 49(1-2): 35-36.
- Saini, A., S.K. Reddy and N. Jawali, 2008. Intra-individual and intra-species heterogeneity in nuclear rDNA ITS region of *Vigna* species from subgenus *Ceratotropis*. *Genet. Res.*, 90: 299-316.
- Salina, E.A., K.Y. Lim, E.D. Badaeva, A.B. Shcherban, I.G. Adonia, A.V. Amosova, T.E. Samatadze, T.Y. Vatolina, S.A. Zoshchuk and A.R. Leitch, 2006. Phylogenetic reconstruction of *Aegilops* section and the evolution of tandem repeats in the diploid and derived wheat polyploids. *Genome*, 49(8): 1023-1035.
- Sharma, S., S. Rustgi, H.S. Balyan and P.K. Gupta, 2002. Internal Transcribed Spacer (ITS) sequences of ribosomal DNA of wild barley and their comparison with ITS sequences in common wheat. *Barley Genet. Newslett.*, 32: 38-45.

- Singh, A., R.M. Devarumath, S. Ramarao, V.P. Singh and S.N. Raina, 2008. Assessment of genetic diversity and phylogenetic relationships based on ribosomal DNA repeat unit length variation and Internal Transcribed Spacer (ITS) sequences in chickpea (*Cicer arietinum*) cultivars and its wild species. *Genet. Resour. Crop. Evol.*, 55: 65-79.
- Sliai, A.M. and S.A.M. Amer, 2011. Contribution of chloroplast DNA in the biodiversity of some *Aegilops* species. *Afr. J. Biotechnol.*, 10(12): 2212-2215.
- Suh, Y., L.B. Thien, H.E. Reeve and E.A. Zimmer, 1993. Molecular evolution and phylogenetic implications of ribosomal DNA in Winteraceae. *Am. J. Bot.*, 80: 1042-1055.
- Sun, Y., D.Z. Skinner, G.H. Liang and S.H. Hulbert, 1994. Phylogenetic analysis of sorghum and related taxa using internal transcribed spacers of nuclear ribosomal DNA. *Theor. Appl. Genet.*, 89: 26-32.
- Takumi, S., S. Nasuda, Y.G. Liu and K. Tsunewaki, 1993. Wheat phylogeny determined by RFLP analysis of nuclear DNA. 1. Einkorn wheat. *Jpn. J. Genet.*, 68: 73-79.
- Terachi, T., Y. Ogihara and K. Tsunewaki, 1984. The molecular basis of genetic diversity among cytoplasm of *Triticum* and *Aegilops*. III. Chloroplast genomes of the M and modified M genome-carrying species. *Genetics*, 108: 681-695.
- Terachi, T., Y. Ogihara and K. Tsunewaki, 1988. The *rbcl* genes in wheat and several *Aegilops* species with divergent chloroplast genomes. *Proceeding of the 7th International Wheat Genetics Symposium*. Cambridge, UK, pp: 789-795.
- Torres, R.A., M. Ganal and V. Hemleben, 1990. GC balance in the internal transcribed spacers ITS 1 and ITS 2 of nuclear ribosomal RNA genes. *J. Mol. Evol.*, 30: 170-181.
- Tsunewaki, K. and Y. Ogihara, 1983. The molecular basis of genetic diversity among cytoplasm of *Triticum* and *Aegilops*. 11. On the origin of polyploid wheat cytoplasm as suggested by chloroplast DNA restriction fragment patterns. *Genetics*, 104: 155-171.
- Van Slageren, M.W., 1994. Wild wheats: A monograph of *Aegilops* L. and *Amblyopyrum* (Jaub and Spach) Eig. (Poaceae): A Revision of All Taxa Closely Related to Wheat, Excluding Wild *Triticum* Species, with Notes on Other Genera in the Tribe Triticeae, Especially *Triticum*. Agricultural University, Wageningen.
- Wan, Y., D. Wang, P.R. Shewry and N.G. Halford, 2002. Isolation and characterization of five novel high molecular weight subunit genes from *Triticum timopheevi* and *Aegilops cylindrica*. *TAG*, 104(5): 828-839.
- Wang, G.Z., N.T. Miyashita and K. Tsunewaki, 1997. Plasmon analysis of *Triticum* (wheat) and *Aegilops*: PCR-single-strand conformational polymorphism (PCR-SSCP) analyses of organellar DNAs. *Proc. Natl. Acad. Sci. USA*, 94: 14570-14577.
- Wang, G.Z., Y. Matsuoka and K. Tsunewaki, 2000. Evolutionary features of chondriome divergence in *Triticum* (wheat) and *Aegilops* shown by RLFP analysis of mitochondrial DNAs. *Theor. Appl. Genet.*, 100: 221-231.
- Waters, E. and B. Schaal, 1996. Biased gene conversion is not occurring among rDNA repeats in the Brassica triangle. *Genome*, 39: 150-154.
- Wendel, J., A. Schinabel and T. Seelanan, 1995. Bidirectional interlocus concerted evolution following allopolyploid speciation in cotton (*Gossypium*). *Proc. Natl. Acad. Sci. USA*, 92: 280-284.
- White, T.J., T. Bruns, S. Lee and J. Taylor, 1990. Amplification and Direct Sequencing of Fungal Ribosomal RNA Genes for Phylogenetics, In: Innis, M.A., D.H. Gelfand, J.J. Sninsky and T.J. White (Eds.), *PCR Protocols, a Guide to Methods and Applications*. Academic Press, San Diego, California, USA, pp: 315-322.
- Yen, C., J.L. Yang and Y. Yen, 2005. The modern genetic concept of the genera in the tribe Triticeae (Poaceae). *Acta Phytotax. Sin.*, 43: 82-93.
- Zhang, W., L.J. Qu, H. Gu, W. Gao, M. Liu, J. Chen and Z. Chen, 2002. Studies on the origin and evolution of tetraploid wheats based on the Internal Transcribed Spacer (ITS) sequences of nuclear ribosomal DNA. *Theor. Appl. Genet.*, 104: 1099-1106.
- Zhang, L.Y., C. Ravel, M. Bernard, F. Balfourier, P. Leroy, C. Feuillet and E.P. Sourdill, 2006. Transferable bread wheat EST-SSRs can be useful for phylogenetic studies among the Triticeae species. *Biomed. Life Sci. TAG Theoret. Appl. Genet.*, 3: 407-418.