

## Some Age Related Attributes of *Macrobrachium macrobrachion* (Herklots, 1851) from Luubara Creek in Ogoni Land, Niger Delta, Nigeria

<sup>1</sup>S.N. Deekae and <sup>2</sup>J.F.N. Abowei

<sup>1</sup>Department of Fisheries and Aquatic Environment, Faculty of Agriculture, Rivers State University of Science and Technology, Port Harcourt, Rivers State, Nigeria

<sup>2</sup>Department of Biological Sciences, Faculty of Science, Niger Delta University, Wilberforce Island, Amassoma, Bayelsa State, Nigeria

**Abstract:** Some Age related attributes of *Macrobrachium macrobrachion* from Luubara Creek in Ogoni Land, Niger Delta, Nigeria was studied for a period of two years (January 2006 - December, 2007). The age composition of the shrimps estimated from the length-at-age data obtained from Paul's integrated plots based on the carapace length showed that; in 2006 males of one year plus represented 35.63% of the sample population while two years plus constituted 55.17% of the sample population. Shrimps of three year plus constituted 6.89% of the sample population while of four years plus represented only 2.29% of the sample population. Shrimps less than one year were not observed. In 2007, male shrimps less than one year of age constituted 32.09% of the sample population while those of one year plus constituted 44.44% or the majority of the catch. Shrimps aged two years plus were 16.04% of the sample population and those of three years plus were 7.40% of the sample population. There were no shrimps of more than four years during the study in 2007. In 2006, there were no female shrimps below one year. Shrimps of one year plus constituted 24.77% of the sample population; shrimps of two years plus constituted 62.83% of the sample population while shrimps of three years plus constituted 12.38% of the population. There were no female shrimps of above four years in 2006. In 2007, there were also no shrimps above four years. Female shrimps of one year plus was 31.11% of the sample population while those more than two years were 54.54% of the sample population. Female shrimps of three years plus constituted only 14.28% of the sample population in 2007. In the berried female shrimps, no shrimp was recorded below one year of age in 2006. Berried female shrimp within one year plus constituted 60% of the sample population while those of two years plus constituted 40% of the sample population. Shrimps of three years age were not found. In 2007, no berried females shrimps caught were below one year. Shrimps of two years plus constituted 70% of the sample population. The relative ages and modal lengths (i.e., length - at - age) of *M. macrobrachion* obtained through Pauly's Integrated distribution plots are showed that the ages attained in males ranged from +1 year to 4 years while the female attained a maximum age of 3 years. It was observed that the growth in length was faster in the males than in females. The annual increment of males was between 3-10 mm in 2006, while berried females had between 3-6.5 mm annual increments. Individual growth also decreased according to age. Individual growths between two age groups were considerably low.

**Key words:** Age, *Macrobrachium macrobrachion*, Luubara Creek, Ogoni Land, Niger Delta, Nigeria

### INTRODUCTION

*Macrobrachium macrobrachion* is a fresh water shrimp; belonging to the Phylum, *Arthropoda*; Class, *Crustacea*; Subclass, *Malacostraca*; Series, *Eumalacostraca*; Order, *Decapoda*; Suborder, *Natantia*; Section, *Caridea*; Family, *Palaemonida*; Genus, *Macrobrachium*; Species, *M. macrobrachion* Powell (1980). It can also be found in low salinity brackish water (Powell, 1985).

The body is divided into three main divisions: the head, thorax and abdomen. The head and thorax are joined to form a cephalothorax, which carries the

mandibles, flagella, rostrum and the eyes containing a stalk and has five pairs of walking legs. The abdomen has six body segments with the last segment bearing a uropod or telson. The other five segments bear swimming apparatus known as swimmerets. A definite feature of *Macrobrachium* is that the second walking legs are modified to form the chelae. Most species are distinctively colored having either blue or brownish colors. The legs also have definitive features such as hairs or furs.

Significant differences exist between the male and female. Mature males are considerably larger than females and the second walking leg is much thicker. The

cephalothorax is also proportionally larger in the male than female while abdomen is narrower in the female. The genital pores of the male are between the bases of the fifth walking leg (New and Singholka, 1982). The female's genital pores situate at the base of the third walking legs. The pleura of the abdomen are lower and broader in the female than in the male. The pleura of the female form a brood chamber in which the eggs are carried between laying and hatching. A ripe ovigerous female can easily be identified because the ovaries can be seen as large orange-colored mass occupying a large portion of the cephalothorax.

The gear used for collecting the shrimp is locally known as "Kara". It is cone shaped and has two non-return value mechanisms at the center of the trap. The trap is constructed from either the blades of bamboo plant or blades of raffia fronds which are woven around three round frames made from cane. The total length of each trap was between 0.95 and 1m while the opening aperture was between 25 and 30 cm. Fresh palm oil fruits were used as bait to set the trap along the creek lets against the water current.

Shrimps and prawns of the genus *Macrobrachium* and *Penaeus* are highly cherished by the people of the Niger Delta. They are used as condiments in the preparation of food because of their high protein value (Umoh and Bassir, 1977; Deekae and Idoniboye-Obu, 1995). They are highly priced and are in high demand in the market (Marioghae, 1990). It has been observed that there is significant reduction of the natural stock of shrimps in our coastal waters (Nwosu, 2007). This may be due to environmental degradation which is detrimental to the abundance and life cycle of *M. macrobrachion*. Also, there are few fishers now to exploit the available species as a result of rural migration.

The unfriendly fishing methods of local fishers who use poisons and chemicals are affecting the shrimp catch. Therefore understanding the biology, environmental parameters and population structure is essential to optimize production from the wild. The shrimp *M. macrobrachion* is exploited in Luubara creek Rivers State in large quantities yet there are no reports on the population biology of this species in the area. A study of the mortality, exploitation of *Macrobrachium macrobrachion* from Luubara creek provides base line data for management decision in the management of the species in the area and similar water bodies.

Age and growth information provides a lot of tools that are used in fishery management. Carlendar (1955) noted that the data on age of a fish or shrimp can provide the following tools in fishery management:

- The general background information needed for management decisions.
- It aids in diagnosis of management needs such as the recognition of overcrowding and stunting.

- It provides information for evaluating the effects of various environmental or hereditary factors by comparing growth rates and year - class abundance.
- It provides information to evaluate the effect of abundance
- Age data can give growth and mortality rates which can be used to estimate optimum yields and possible effects of catch regulations.
- Age and growth data can be used to predict year -class abundance.
- Production rates can be determined with estimates of the age and growth data.

In the temperate regions growth can be estimated by counting of the growth zones or annual marks which appear on the hard parts of fishes or shrimps as a result of definite climate seasons. This is not the case in the tropical regions where climatic seasons are not clearly defined. Some methods have evolved and are used for the estimation of age in fishes and shrimps. According to Bhukaswan (1980) the methods of age and growth determination are:

- Interpretation of and counting of growth zones or annual marks, which appear in the hard parts of fishes.
- Marking or tagging and recapture
- Comparison of length frequency distribution.

The age of fishes can be determined by the examination of hard parts of the fish such as scales, otoliths, spine and finrays, opercular bones and vertebrae. The examination of annual marks on scale is a popular method used in aging fish (Fagade, 1974; Krupka, 1974; Bastl, 1974; Beamish and Fournier, 1981; Rao and Tapia, 1999). Many authors have reported to use otolith as a method to determine age (Itoh and Tsuji, 1996). According to Morales-Nin, (2000) and Megalofonou (2006), the aging of fish through the use of hard parts poses some problems for fisheries scientists in the tropics because there are no definite seasons which can show growth differences. In the temperate regions age and growth can be estimated by counting of growth zones or marks of the fish as a result of definite climatic seasons. This is not the case in the tropical regions where the climatic conditions are fairly constant. Thus aging fish based on the hard parts has some limitations in the tropics.

Marking, tagging and subsequent recapture has been reported to be used to determine age and growth in fish, shrimps and other crustacean (King, 1995, Siddek, 1990). The method can be used on large size fishes but where the fish is small, marking or tagging may become a problem. It may cause mortality of fish and the fish may lose their marks. When the marks are lost fishermen may not be able to recognize the fish (or shrimp) and the method may not be useful.

Data of length frequency distribution of fishes and shrimps can be used to determine age. This method was first introduced by Peterson (1892) as reported by Sparre *et al.* (1989). It is based on taking the length data of a shrimp or fish for a period, usually about one year and pooling the data together. From the data, particular cohorts or different age groups can be determined when plotted on a graph. The length frequency method has undergone some modifications by being combined with the modal class progression analysis to produce the integrated length frequency method (Pauly, 1983; Jones, 1984). This method has been used to estimate the age of shrimps (Garcia and Reste, 1981; Garcia, 1985; Enin, 1995; Enin *et al.*, 1995; Mohammed, 1979) and fish (Nawa, 1985; King, 1996a, b; Ezekiel, 2001; Abowei and Hart, 2007).

One advantage of the length - frequency method is that, it takes into account the influence of the whole spawning season and the production of several broods per annum, a characteristic of fishes and shrimps. Use of length frequency data has become a very important tool in aging of shrimps and fish in the tropics where growth marks and seasonal differences are not properly defined. It should be noted that it is only when samples are large that unbiased parameters can be obtained. Methods of obtaining data from length frequency distribution include the Bhattacharya's method; modal progression analysis; the probability paper and parabola method and computer-based frequency analysis.

This method involves producing a length frequency distribution graph of the sample. The Bhattacharya method basically separate normal distributions, each representing a 'cohort' of fish or shrimp from the overall distribution. The age of each group or 'cohort' can then be estimated by using mathematical formulae such as regression analysis (Sparre *et al.*, 1989).

Recently, computer programmes have been developed incorporating this method. Such packages include the (Length Based Fish Stock Assessment LSFA (Sparre, 1987) and the complete Electronic Length Frequency Analysis ELEFAN (Gayanilo *et al.*, 1988).

The age of shrimps can be estimated by modal progression analysis. It is a modification of the Bhattacharya method. The mean lengths of the cohorts are determined from the Bhattacharya plots using the Von Bertalanffy formulae and regression analysis.

Sparre *et al.* (1989) observed that modal progression analysis is suitable for short live species such as shrimps which do not have many 'cohorts' or age groups in the growth pattern.

This method was developed by Cassie (1954) as reported by Sparre *et al.* (1989). This method is based on the fact that when a normal distribution is plotted on a probability paper, the resultant curve becomes linear. It is easy to spot cumulated distribution on such a line. In the case of the parabola method the normal distribution is

transformed by taking natural logarithms of the length frequency data. The parabolas are then fitted to the log-transformed numbers of the length frequency data.

Computer packages have recently been developed to determine the ages and other growth parameters of fishes and shrimp. These programmes include the complete Electronic Frequency Analysis (ELEFAN) written in the computer language BASIC (Gayanilo *et al.*, 1988). It deals with the estimation of growth parameters using length frequency analysis. It is a revised version of ELEFAN I developed by Pauly and David (1981). There is also the Normal Separation of fish-length frequencies in distributed components (NORMSEP) programme developed by Hasselbald and Tomlinson (1971) which has been extended by Schnute and Fournier (1980) to include estimation of growth parameters and by Sparre (1987) to deal with time series and a seasonalized Von Bertalanffy growth curve. Sparre (1987) also introduced the Length Based Fish Stock Assessment (LSFA) programme in the Bhattic programmed and is also being used in length frequency analysis. There is also the FAO ICLARM Stock Assessment Tool (FiSAT) (Gayanilo and Pauly, 1997). It combines LSFA and complete ELEFAN into one package. It is now the recognized programme for fish stock assessment.

These computer programmes are being revised from time to time to ensure adaptation, efficiency and easy usage. Nevertheless, Sparre *et al.* (1989) has noted that computer based frequency analysis has some limitations. In computer analysis, the data is usually coded and when a large pool of data is used mistakes could be made when coding. Most population parameters can be generated after initial input has been made Sparre *et al.* (1989) observed that the researcher may be biased in selecting the population parameters as given by the computer. Another major problem of computer analysis is to resolve mixed distribution of fishes especially for tropical species which are multi-species. Separation into the various components is usually difficult and time consuming.

Growth studies are very important tools in fishery management. It is the growth of individuals in the population that determines the amount of stock available for the fishery (King, 1995). Growth studies provide information for the evaluation of production; stock sizes; recruitment; mortalities and the status of a shrimp population (Ling, 1969). The growth process is specific for each species of shrimp or fish.

Many factors affect the growth of aquatic organisms and include; food supply; parasitism; competition; predation and environmental factors (salinity; pH; rainfall). Others are presence of adequate anions and cations in the water body; amount of dissolved gases (especially oxygen) and the quantity of pollutants in water. Age and growth studies always go together since age is determined by growth.

Growth may be described as indeterminate for those organisms that have capacity for sustained though diminishing growth throughout their lives if sufficient food is available. Members of such species may be of variable sizes at the same ages in contrast to fishes or shrimp which has determinate growth having more definite sizes at any one age. In other words, fishes and shrimps in contrast to birds and mammals of the terrestrial vertebrate groups do not cease growth after attainment of sexual maturity. They continue their growth though only at slow rates.

**Types of growth:**

**There are three types of growth:**

**Absolute growth:** This is considered as the average size of fish (or shrimp) at each age and is estimated as:

Abs. Growth =  $TL_n + 1 - TL$  (Ehrhardt, *et. al.*, 1983) (1)  
 Where  $TL_n + 1$  = Total length attained with new age  
 $TL$  = Original total length

**Relative growth:** This refers to the increment between two age groups divided by the length at the younger age.

$$h = \frac{L_2 - L_1}{L_1} \quad (\text{Ehrhardt } et \text{ al.}, 1983) \quad (2)$$

where;  $h$  = Relative growth  
 $L_1$  = Length-at-age one  
 $L_2$  = Length-at-age two

**Instantaneous Rate of Growth:** It is also called growth co-efficient. It refers to the difference between the natural logarithms (ln) of length (or weight) for consecutive age groups.

$$g = L_n L_2 - L_n L_1 \text{ or } g = L_n W_2 - L_n W_1 \quad (\text{Ehrhardt } et \text{ al.}, 1983) \quad (3)$$

where;  $g$  = Growth co-efficient  
 $L_n L_1$  = Logarithm of original length  
 $L_n L_2$  = Logarithm of final length  
 $L_n W_1$  = Logarithm of original weight  
 $L_n W_2$  = Logarithm of final weight

Length is the more frequently measured attribute of growth perhaps due to ease of measurement and the relationship that has been developed between length and weight.

Growth parameters are generally estimated using mathematical models (Ricker, 1975; Gulland, 1988). These models are usually based on certain assumptions. These include.

- The models can give or predict the size in terms of length or weight at any given age which agrees with the observed data of size and age.
- The model should be able to make predictions close to reality.

- The model (mathematical form) must be such that it can be incorporated readily into existing models.

According to Sparre *et al.* (1989), the most popular growth equation is the Von Bertalanffy (1938) growth equation, VGBF, expressed as:

$$L(t) = [L_{\infty} 1 - \exp(-k(t - t_0))] \quad (4)$$

where;

$L_{\infty}$  = Asymptotic length  
 $k$  = Growth co-efficient  
 $t_0$  = Hypothetical age at zero length  
 $L_{(t)}$  = Length at age (t)  
 $\exp$  = Constant

The importance of this model is that different growth curves can be created for each set of the growth parameters  $L_{\infty}$ ,  $k$  and  $t_0$ . It is therefore possible to use the model to describe the growth pattern of both finfish and shellfish including shrimps. Growth parameters differ from species to species and from stock to stock and even differ within the same stock (Sparre *et al.*, 1989). In calculating the growth parameters of a stock of fish or shrimp, length frequency data are generally used. For example, data on length - at - age ( $l_t$ ) are required to fit the three parameters -  $L_{\infty}$ ,  $k$  and  $t_0$ . Computer programmes now exist for solving the Von Bertalanffy growth parameters for better results.

This method was reported by Sparre *et al.* (1989). It is based on the original Von Bertalanffy growth equation. It was noted that if fish length at one age was plotted against the length of next younger age, the result was a straight line for many populations. He observed that if the slope of the plot was less than one, a Von Bertalanffy growth curve can be fitted to the data. The parameters  $L_{\infty}$  and  $k$  are estimated from the Ford - Walford plot essentially consists of a rewritten version of VBGF as follows:

$$L_{t+1} = a + b l_t \quad (5)$$

where;

$a$  = regression constant (the intercept)  
 $b$  = co-efficient  
 $l_t$  = length at age (t)  
 $L_{t+1}$  = Length at the next age

It is important to note that  $l_t$  and  $l_{t+1}$  must be separated by a constant time interval.

$$L_{\infty} = \frac{a}{1 - b} \quad (6)$$

The slope of a Walford line is equal to  $e^{-k}$ , thus the natural log of the Walford slope with the sign changed

provides an estimate of growth co-efficient  $k$  written as:  $K = \log^c$ .

The values of  $t_0$  and  $L_t$  may be estimated from the growth equation after estimates of  $K$  and  $L_\infty$  have been made or a mathematical procedure may be used to solve for the three parameters simultaneously.

This method was suggested by Wetherall (1986) for the estimation of asymptotic length,  $L_\infty$  and ratio of total mortality and growth co-efficient,  $Z/K$ . In using this method a large length frequency data is needed as a representative of a steady population usually by pooling samples obtained over many months. The length frequency data are then broken into several size classes. The lowest limit of each class size acts as the cut-off length ( $L^1$ ). The mean length ( $\bar{L}$ ) of all fishes or shrimp from each  $L^1$  upward is computed. A series of pairs of  $L^1$  and  $\bar{L}$  are obtained. The difference between each pair of  $L^1$  and  $\bar{L}$  ( $\bar{L} - L^1$ ) is plotted against the corresponding  $L^1$  and a regression line is fitted to that part of the curve pertaining to fish (shrimp) fully recruited to the gear. When the regression line is fitted the point that the line cuts the  $L^1$  gives the estimate of  $L_\infty$ . It also gives an estimate of  $Z/K$  which is given as:

$$Z / K = \left( \frac{1+b}{-b} \right) \quad (7)$$

The Wetherall method has been modified by Pauly (1986) and is now incorporated into the FiSAT computer programme.

Some Age related attributes of *Macrobrachium macrobrachion* from Luubara Creek in Ogoni Land, Niger Delta, Nigeria provides a lot of tools that are used in fishery management: General background information needed for management decisions; aids in diagnosis of management needs such as the recognition of overcrowding and stunting; information for evaluating the effects of various environmental or hereditary factors by comparing growth rates and year - class abundance; information to evaluate the effect of abundance growth and mortality rates which can be used to estimate optimum yields and possible effects of catch regulations; predict year - class abundance and production rates of the fishery and similar water bodies.

## MATERIALS AND METHODS

**Study area:** The study was carried out in Luubara Creek in Ogoni Land, Niger Delta, Nigeria was studied for a period of two years (January 2006 - December, 2007). The creek is a tributary of the Imo River and is located between Longitudes  $7^{\circ}15' - 7^{\circ}32'E$  and Latitudes  $4^{\circ}32' - 4^{\circ}37'N$  in the Eastern part of the Niger Delta. The upper

part of the creek extends from Bori and meanders through Wiyiakaara, Luegbo, Duburo and joins the Imo River at Kalooko.

The creek is divided into two distinct sections brackish water and freshwater. The brackish water stretch is between Bane and Kalooko while the freshwater stretch extends from Bane to Bori. The brackish water area has the normal mangrove vegetation comprising of trees such as *Rhizophora racemosa*, *Aveenia africana*, *Laguncularia racemosa* etc., whereas the freshwater has dense vegetation comprising of large trees, various palms and aquatic macrophytes at the low intertidal zone. In freshwater area are *Cocos* species, *Eliasis* species, *Nymphaea* species, *Lemna* species and *Raffia* species.

It is characterized by high ambient temperature usually about  $25.5^{\circ}C$  and above; high relative humidity which fluctuates between 60 and 95% and high rainfall averaging about 2500 mm (Gibo, 1988). This high rainfall often increases the volume of water in the creek hence providing good fishing opportunity for the residents. Fishing is one of the major activities going on along the creek because it is the main water route of the Khana people in Ogoni area of the Niger Delta.

The fishes caught in the area include *Chrysichthys auratus*, *C. nigrodigitatus*, *Hydrocynus forskalii*, *Clarias gariepinus*, *Pellonula leonensis*, *Malapterurus electricus*, *Gymnarchus niloticus*, *Synodontis nigri* *Hepsetus odoe*, *Hernichromis fasciatus*, *Tilapia zilli*, *Tilapia guineensis*; *Sarotherodon melanotheron* and *Eleotris senegalensis* and shellfish (crabs and shrimps) especially *Uca tangeri* *Callinectes amnicola*, *Goniopsis pelli*, *Cardisoma armatum* *M. macrobrachion*, *M. vollenhoveni*, *M. equidens*, *Palaemonetes africanus*, *Caridina africana* and *Desmocarais tripisnosa*.

**Specimen sampling:** The shrimp samples were collected fortnightly from three stations along the creek: namely Wiyiakaara, Luegbo and Duburo. Selection of the stations was purposefully based on fishing activities, ecological zonation and accessibility of site. For each station five fishermen were engaged and three traps were used. At each station the fisher men set the three sets of traps against the water current among aquatic macrophytes and left them overnight. The traps were retrieved the following day after about twelve hours corresponding to another low tide. The shrimps collected at each station were sorted into male and female; females were later separated into berried (ovigerous) and non-berried (non-ovigerous). Sampling lasted for twenty-three months from January 2006 to November 2007. The shrimp samples were then preserved in 4% formaldehyde and transported to the RSUST Fisheries laboratory for analysis after each day's sampling. The species was identified by use of the keys of Powell (1980; 1982) and Holthius (1980).

For each shrimp the Total length (the distance from the tip of the rostrum to the end of telson) and the

carapace length (the distance from the base of rostrum to the first body segment) was measured with a Vernier caliper to the nearest 0.1 mm. The shrimps were then weighed with an Ohaus balance to the nearest 0.1 g. Measurements were taken for each monthly collection and recorded accordingly.

The age structure of the shrimps obtained during the period of study was estimated using the modified Peterson method also known as the Integrated method (Pauly, 1983; Sparre *et al.*, 1989). Monthly length measurements were pooled and plotted on a graph to obtain length frequency distribution for the period of study at 4mm class interval. The same process was repeated along the time axis on the assumption that growth pattern repeats itself from year to year. Pauly's Integrated frequency model is based on the assumption that growth in length in shrimps is rapid initially, then increasing smoothly, passing through most of the discrete peaks. The annual growth pattern can be obtained from the Von Bertalanffy growth formula (Sparre *et al.*, 1989) as follows:

$$L_t = L_{\infty} (1 - \exp [-K (t - t_0)])$$

where;

- $L_t$  = length - at - age
- $t_0$  = is a constant and represents length - at - age zero
- $L_{\infty}$  = representing maximum possible length
- $K$  = growth co-efficient

$L_{\infty}$  and  $K$  were obtained from the Von Bertalanffy plots (Pauly, 1983; Sparre *et al.*, 1989; King, 1995) using the formular:

$$L_t + 1 = a + bL_t \tag{8}$$

where;

- $a$  = axis intercept
- $b$  = slope of regression line
- $L_t$  = Length- at- age  $t$
- $L_t + 1$  = Length - at- next age

$L_t$  and  $L_t + 1$  are separated by a constant time interval of one year.

Hence:

$$L_{\infty} = \frac{a}{1-b} \text{ and } K = -\log b \tag{9}$$

Estimate of  $t_0$  was made by plotting:

$$t_0 - t_0 = \left[ \frac{L_{\infty} - L_t}{L_{\infty}} \right] = a + bt \tag{10}$$

$$t_0 = \left[ \frac{L_{\infty} - L_t}{L_{\infty}} \right] \tag{11}$$

- $a$  = axis intercept
- $b$  = Slope of regression line

This method is incorporated in the FiSAT programme and the values of  $L_{\infty}$  and  $K$  obtained.  $L_{\infty}$  was also estimated by using the method of Wetherall (1986) as modified by Pauly (1986). In Wetherall method the monthly length frequency data of the shrimps were pooled together. The length frequency pattern was then broken into different size classes. The lowest limit of each size class acts as the cut off length ( $L^1$ ). The mean length ( $\bar{L}$ ) of all shrimp from each cut off length ( $L^1$ ) upwards is computed. A series of pairs of  $L^1$  and  $\bar{L}$  were obtained. The difference between each pair of  $L^1$  and  $\bar{L}$  ( $\bar{L} - L^1$ ) was plotted against the corresponding  $L^1$  and a regression line fitted to that part of curve pertaining to shrimp fully recruited to the gear. The regression line has the form  $\bar{L} - L^1 = a + b L^1$ . The point where  $L^1$  cuts the regression line gives the estimate  $\frac{Z}{K} = \frac{(1+b)}{-b}$  and  $L_{\infty} = \frac{-a}{-b}$  of  $\bar{L}$  or can be presented as indicated:

The Wetherall method has been incorporated in the FiSAT computer programme (Gayanilo and Pauly, 1997)

hence, used to obtain  $L^1$  and  $t$ , and  $\frac{Z}{K}$

## RESULTS AND DISCUSSION

The age composition of the shrimps was estimated from the length-at-age data obtained from Paul's integrated plots based on the carapace length and shown in Table 1. In 2006 males of one year plus represented 35.63% of the sample population while two years plus constituted 55.17% of the sample population. Shrimps of three year plus constituted 6.89% of the sample population while of four years plus represented only 2.29% of the sample population. Shrimps less than one year were not observed (Table 1a). In 2007, male shrimps less than one year of age constituted 32.09% of the sample population while those of one year plus constituted 44.44% or the majority of the catch. Shrimps aged two years plus were 16.04% of the sample population and those of three years plus were 7.40% of the sample population. There were no shrimps of more than four years during the study in 2007 (Table 1b).

In 2006, there were no female shrimps below one year. Shrimps of one year plus constituted 24.77% of the

Table 1: Age composition of *M. Macrobrachion* in Luubara creek

Age	Modal length (mm)	No. of Shrimps	Age Composition
<b>a Male (2006)</b>			
0	3	0	0%
1 <sup>+</sup>	11	31	35.63%
2 <sup>+</sup>	14	48	55.17%
3 <sup>+</sup>	24	6	6.89%
4 <sup>+</sup>	28	2	2.29%
<b>b Male (2007)</b>			
0	6.5	26	32.09%
1 <sup>+</sup>	14.0	36	44.44%
2 <sup>+</sup>	21.5	13	16.04%
3 <sup>+</sup>	24.5	6	7.40%
4 <sup>+</sup>	27.0	0	0%
<b>c Female (2006)</b>			
0	2	0	0
1 <sup>+</sup>	6	28	24.77
2 <sup>+</sup>	12	71	62.83
3 <sup>+</sup>	17	14	12.38
<b>d Female (2007)</b>			
0	2	0	0
1 <sup>+</sup>	8	24	31.11
2 <sup>+</sup>	13.5	42	54.54
3 <sup>+</sup>	19.0	11	14.28
<b>e Berried female (2006)</b>			
0	3	0	0
1 <sup>+</sup>	8.5	57	60
2 <sup>+</sup>	15	38	40
3 <sup>+</sup>	20.5	0	0
<b>f Berried female (2007)</b>			
0	3.2	0	0
1 <sup>+</sup>	10.4	35	70
2 <sup>+</sup>	17.0	15	30
3 <sup>+</sup>	21.5	0	0

sample population; shrimps of two years plus constituted 62.83% of the sample population while shrimps of three years plus constituted 12.38% of the population.

There were no female shrimps of above four years in 2006 (Table 1c). In 2007, there were also no shrimps above four years. Female shrimps of one year plus was 31.11% of the sample population while those more than two years were 54.54% of the sample population. Female shrimps of three years plus constituted only 14.28% of the sample population in 2007 (Table 1d).

In the berried female shrimps, no shrimp was recorded below one year of age in 2006. Berried female shrimp within one year plus constituted 60% of the sample population while those of two years plus constituted 40% of the sample population (Table 1e). Shrimps of three years age were not found. In 2007, no berried females shrimps caught were below one year. Shrimps of two years plus constituted 70% of the sample population (Table 1f).

The relative ages and modal lengths (i.e., length - at - age) of *M. macrobrachion* obtained through Pauly's Integrated distribution plots are shown in Table 2 and 3. The ages attained in males ranged from +1 year to 4 years while the female attained a maximum age of 3 years. It was observed that the growth in length was faster in the males than in females.

The annual increment of males was between 3-10 mm in 2006, while berried females had between 3-6.5 mm

annual increment. Individual growth also decreased according to age. Individual growths between two age groups were considerably low.

The result of Pauly's Integrated Length Frequency for relative length - at - age of *M. macrobrachion* demonstrated a population with 3-4 cohorts. The shrimps fall within one to three years. Most of the shrimps in the sample population studied were above one year. The ages attained in males ranged from one year to three years while females attained a maximum of three years. Berried females also attained a maximum age of three years. This is similar to the result of Enin (1995) and Enin *et al.* (1995), which suggested a maximum age of two and half years for *M. macrobrachion*.

### CONCLUSION

- The relative length - at - age of *M. macrobrachion* from demonstrated a population with 3-4 cohorts.
- The shrimps fall within one to three years.
- Most of the shrimps in the sample population studied were above one year. The ages attained in males ranged from one year to three years while females attained a maximum of three years.
- Berried females also attained a maximum age of three years.
- The results confirmed other studies of the species from similar water bodies.

Table 2: Length-at-age and annual length increment of the various sexes of *Macrobrachium macrobrachion* in Luubara creek (2006-2007)

Age	Modal length (mm)	Annual Increment	Length + 1
<b>Male 2006</b>			
0+	3	3	11
1+	11	8	14
2+	14	3	24
3+	24	10	28
4+	28	4	0
<b>Female 2006</b>			
0+	2	2	6
1+	6	4	12
2+	12	6	17
3+	17	5	0
4+	0	0	0
<b>Berried female 2006</b>			
0+	3	3	8.5
1+	8.5	5.5	15
2+	15	6.5	20
3+	20.5	5.5	0
4+	0	0	0
<b>Male 2007</b>			
0+	6.5	6.5	14.0
1+	14.0	7.5	21.5
2+	21.5	7.0	24.5
3+	24.5	3.0	27.0
4+	27.0	2.5	0
<b>Female (2007)</b>			
0+	2	2	8
1+	8.0	6.0	13.5
2+	13.5	5.5	19.5
3+	19.5	6.0	0
<b>Berried Female (2007)</b>			
0+	3.2	3.2	10.43
1+	10.4	7.2	17.0
2+	17.0	6.6	21.0
3+	21.5	4.5	0
4+	0	0	0

Table 3: Length-at-age and absolute increment in male, female and berried female *M. Macrobrachion* for Luubara creek (2006 - 2007)

Age (Yrs)	Modal Length (CL) mm	Absolute Increment (m)	$L_{t+1}$ (mm)
<b>2006</b>			
Male 1 <sup>+</sup>	11	3	14
2 <sup>+</sup>	14	10	24
3 <sup>+</sup>	24	4	28
4 <sup>+</sup>	28	-	-
Female 1 <sup>+</sup>	6	6	12
2 <sup>+</sup>	12	5	17
3 <sup>+</sup>	17	-	-
B female 1 <sup>+</sup>	8.5	6.5	15
2 <sup>+</sup>	15	5.5	20.5
3 <sup>+</sup>	20.5	-	-
<b>2007</b>			
Male 1 <sup>+</sup>	14.0	7.5	21.5
2 <sup>+</sup>	21.5	3.0	24.5
3 <sup>+</sup>	24.5	3.5	27.0
4 <sup>+</sup>	27.0	-	-
Female 1	8.0	5.5	13.5
2	13.5	6.0	19.5
3	19.5	-	-
B female 1	3.2	7.2	10.4
2	10.4	7.1	17.5
4	17.5	-	-

**ACKNOWLEDGMENT**

We are grateful to God Almighty for the strength, knowledge and wisdom in researching and writing this

article. We also thank our numerous colleagues, friends, wives and students whose names are too numerous to mention here due to space but contributed immensely to the success of this work.

## REFERENCES

- Abowei, J.F.N. and A.I. Hart, 2007. A study of length-weight relationship, condition factor and age of ten fish species from the lower Nun River, Niger Delta. *Afr. J. Appl. Zool. Environ. Biol.*, 9: 13-19.
- Bastl, I., 1974. The red-breasted bream *Tilapia rendalli* (Boulenger, 1896). *Monograph Biol.*, 24: 311-318.
- Bhukaswan, T., 1980. Management of Asian Reservoir Fisheries. *FAO Fish. Tech. Pap.*, 207: 69.
- Beamish, R.J. and D.A. Fournier, 1981. A method of comparing the precision of a set of age determinations. *Can. J. Fish. Aqua. Sci.*, 38: 982-983.
- Carlander, K.D., 1955. The Standing crop of fish in lakes. *J. Fish. Res. Board Canada*, 12(4): 543-570.
- Cassie, R.M., 1954. Some uses of probability paper in the analysis of size - frequency distributions. *Aust. J. Mar. Fresh Water Res.*, 5: 513-522.
- Deekae, S.N. and T.I.E. Idoniboye-Obu, 1995. Some aspects of commercially important molluscs and crabs in the Niger Delta, Nigeria. *Environ. Ecol.*, 13(1): 136-142.
- Ehrhardt, N.M., P.S. Jacquemin, G.G. Francisco, G.D. German, M.L.B. Juan, O.O. Juan and S.N. Austin, 1983. On the Fishery and Biology of the Giant Squid, *Dosidicus Oloas* in the Gulf of California, Mexico. In: Caddy, J.R. (Ed.), *Advances in Assessment of World Cephalopod Resources*. *FAO Fisheries Technical Paper*, 231: 306-339.
- Enin, U.I., 1995. First estimates of growth, mortality and recruitment parameters of *Macrobrachium macrobrachion* Herklots, 1851 in the Cross River Estuary, Nigeria. *Dana*, 2(1): 29-38.
- Enin, U.I., F. Lowenberg and T. Kunzel, 1995. Population dynamics of the estuarine prawn (*Nematopalaemon hastatus*, Aurivilius 1898) of the Southeast Coast of Nigeria. *Fish. Res.*, 26: 17-35.
- Ezekiel, E.N., 2001. Comparative studies of the flood plains and major rivers in Odhiokwu Ekpeye Niger Delta. *M.Sc. Thesis, University of Port Harcourt, Choba*, pp: 77.
- Fagade, S.O., 1974. Age Determination in *Tilapia melanotheron* (Ruppell) in the Lagos Lagoon, Lagos, Nigeria. In: Bagenal, T.B. (Ed.), *Proceedings of an International Symposium on the Ageing of Fish*. The Gresham Press, Unwin Brother Ltd., Old Working Surrey, pp: 71-77.
- Garcia, S. and L. Reste, 1981. Life cycles, dynamics, exploitation and management of coastal penaeid shrimp stocks. *FAO Fish. Tech. Pap.*, 203: 215.
- Garcia, S., 1985. Reproduction, Stock Assessment Models and Population Parameters in Exploited. In: Rothlisberg, P.C., B.J. Hill and D. Staples (Eds.), *Second Australian National Prawn Seminar*, Cleveland, Queensland Australia, NPS2, pp: 139-158.
- Gayanilo, F.C., M. Soriano and D. Pauly, 1988. A draft guide to the COMPLEAT ELEFAN. *ICLARM Software Project 2*, pp: 65.
- Gayanilo, F.C. and D. Pauly, 1997. *FAO-ICLARM Stock Assessment Tools (FiSAT)*. *FAO Computerised Information Series (fisheries) No. 8*, Rome, pp: 262.
- Gibo, A.E., 1988. Relationship between rainfall trends and flooding in the Niger- Benue River basin. *J. Meteorol.*, 13: 132-133.
- Gulland, J.A., 1988. *Fish Population Dynamics: The Implications for Management*. 2nd Edn., John Wiley and Sons Ltd., Chichester, pp: 422.
- Hasselbald, V. and P.K. Tomlinson, 1971. *NORMSEP. Normal Distribution Separator*. In: Abramson, N.J. (Ed.), *Computer Programs for Fish Stock Assessment*. *FAO Fisheries Technical Paper 101*: 11(1): 1-10.
- Holthius, L.B., 1980. *Shrimps and prawns of the world: An annotated catalogue of species of interest to fisheries*. *FAO Fisheries Synopsis*.
- Itoh, T. and S. Tsuji, 1996. Age and growth of juvenile Southern blue fin tuna *Thunnus maccoyii* based on otolith microstructure. *Fish. Sci.*, 62: 892-896.
- Jones, R., 1984. Assessing the effects of changes in exploitation pattern using length composition data (with notes on VPA and cohort analysis). *FAO Fish. Tech. Pap.*, 256: 118.
- King, R.P., 1995. *Fisheries Biology, Assessment and Management*. *Fishing News Books, Surrey*, pp: 336.
- King, R.P., 1996a. Length - weight relationships of some Nigeria freshwater fishes. *NAGA, ICLARM Q.*, 19(3): 49-52.
- King, R.P., 1996b. Length-weight relationships of Nigerian coastal water fishes. *NAGA, ICLARM Q.*, 19(4): 53-55.
- Krupka, I., 1974. The kurper bream, *Sarotherodon mossambicus* (Trewavas, 1966). *Monograph. Biol.*, 24: 343-349.
- Ling, S.W., 1969. The general biology and development of *Macrobrachium rosenbergii*. *FAO Fish Report (57)*, 3: 589-606.
- Marioghae, I.E., 1990. Studies of fishing methods, gear and marketing of *Macrobrachium* in the Lagos Area. *Nigerian Institute of Oceanography and Marine Research Technical Paper*, No. 53, pp: 20.
- Megalofonou, P., 2006. Comparison of otolith growth and morphology with somatic growth and age in Blue fin tuna. *J. Fish Biol.*, 68: 1867-1878.
- Mohammed, K.H., 1979. Mark - recapture experiments on the Gulf shrimp *Penaeus semisulcatus*, De, Haan, in Kuwait waters. *Shrimp stock valuation and management project, Kuwait*. *Report TF/KUW -6/6/R-10*, pp: 59 (Mimeo).
- Morales-Nin, B., 2000. Review of the growth regulation process of otolith daily investment formation. *Fish. Res.*, 46: 53-67.

- Nawa, I.G., 1985. A study on the growth of *Pseudotolithus elongatus* and *Chrsichthys nigrodigitatus* occurring in the Cross River estuary. Proceedings of the 4th Annual Conference of the Fisheries Society of Nigeria, pp: 162-166.
- New, M.B. and S. Singholka, 1982. Freshwater prawn farming: Manual for the culture of *Macrobrachium rosenbergii*. FAO Fisheries Technical. Paper, 225, pp: 116.
- Nwosu, F., 2007. The Problem of By Catch Associated with Industrial Shrimping: Implications for Inshore Demersal Fisheries in the Niger Delta. In: Zabbey, (Ed.), Small Scale Shrimp Fisheries in Nigeria. Centre for Environment, Human Rights and Rural Development (CEHRD), Eleme, Rivers State CEHRD/TECH/CONSERV/01/2007, pp: 32-48.
- Pauly, D., 1983. Some simple methods for the assessment of tropical fish stocks. FAO Fish. Tech. Pap., 234: 52.
- Pauly, D. and N. David, 1981. ELEFAN, a basic program for the objective extraction of growth parameters from length frequency data. *Mecresforschimg*, 28(4): 205-211.
- Pauly, D., 1986. On improving the operation and use of the ELEFAN programs. Part III: Correcting length - frequency data for effects of gear selection and /or incomplete re-enactment ICLARM. *Fishbyte*, 4(2): 11-13.
- Powell, C.B., 1980. Key to shrimps and prawns (crustacea: decapoda, natantia) of the Niger Delta Basin Development Authority Area. Consultancy Report, pp: 5.
- Powell, C.B., 1982. Fresh and brackishwater shrimps of economic importance in the Niger Delta. Proceeding 2nd Annual Conference Fisheries Society of Nigeria. Calabar, 25-27 January, pp: 254-285.
- Powell, C.B., 1985. The Decapods Crustaceans of the Niger Delta. In: Wilcox, H.B.R. and C.B. Powell, (Eds.), Publication Committee, University of Port Harcourt, pp: 226-238.
- Rao, R. and F. Tapia, 1999. Estimation of size at sexual maturity evaluation of analytical and re-sampling procedures. *Fish. Bull.*, 97: 570-580.
- Ricker, W.E., 1975. Computation and interpretation of biological statistics of fish populations. *Bull. Fish. Res. Board Can.*, 191: 382.
- Schnute, J. and D. Fournier., 1980. A new approach to length - frequency analysis: growth structure. *Can. J. Fish Aqua. Sci.*, 37: 1337-1351.
- Siddek, M.S., 1990. Estimating natural mortality of Kuwait's *Penaeus semisulatus* using a new tag - recapture data analysis method. *ICLARM Fishbyte*, 8(3): 10-12.
- Sparre, P., 1987. Computer programs for fish stock assessment. Length - based fish stock assessment for Apple 11 computers. FAO Fish. Tech. Pap., 101(Suppl. 2): 218.
- Sparre, P., E. Ursin and S.C. Venema, 1989. Introduction to tropical fish stock assessment. Part I. manual FAO Fisheries Technical Paper No. 306. 1 FAO Rome, pp: 337.
- Umoh, I.B. and O. Bassir, 1977. Lesser known sources of protein in some Nigerian peasant diets. *Food Chem.*, 2: 315-329.
- Bertalanffy, L., 1938. A quantitative theory of organic growth. *Hum. Biol.*, 10: 181-243 (cited from Sparre *et al.*, 1989).
- Wetherall, J.A., 1986. A new method for estimating growth and mortality parameters from length - frequency data. *ICLARM Fishbyte*, 4(1): 12-14.