

Stem Biomass of *Anchusa italica* (*Anchusa azurea*) in Shanjan Rangelands, East Azerbaijan, Iran

Ghassem Habibi Bibalani and Hamideh Shadkami-Til

Department of Agriculture, Shabestar Branch, Islamic Azad University, Shabestar, Iran

Abstract: Biomass of plant stem have been studied in this research, because Plants over ground parts can be used for animal grazing, in wind erosion control, to reduce water flow rates, and to increase evaporation and transpiration. In the NW of Iran (East Azerbaijan Province), rangelands previously used to animal grazing were changed to agricultural land use; this vegetation is unsuitable vegetation coverage. We studied *Anchusa italica* (*Anchusa azurea*) to determine its stem biomass characteristics. Data were collected using an accidental sampling methodology (1*1 m). In total, 7 plots were identify and 35 samples were collected for this research. In the minimum, maximum and mean stem biomass for this plant were found to me 18.5, 31.4 and 22 g, respectively.

Key words: *Anchusa italica* (*Anchusa azurea*), Iran, Rangeland, stem biomass

INTRODUCTION

Rangeland ecosystem stabilizing, optimum and continual utilization of the range without studding and knowing the influencing factors on its segments and animal pasturage are of special importance (Bibalani *et al.*, 2011a; Bibalani *et al.*, 2011b; Mozaffarian, 2007; Shadkami-Til and Bibalani, 2010; Shadkami-Til and Bibalani, 2011). There are different methods of evaluating rangelands and all of them have advantages and disadvantages. Factors such as vegetation species composition, annual production, area coverage, plant density, soil surface coverage, constitution, and presence of succulence plants were used (Bidlock *et al.*, 1999; Mogaaddam, 2001) but estimation of these parameters are time consuming and expensive.

Fresquez (Fresquez *et al.*, 1990) reported an increase in vegetative production and forage quality of Blue Grama (Mata-Gonzalez *et al.*, 2002). Benton & Wester (Benton and Wester, 1998) reported an increase in Tobosagrass (*Hilaria mutica*) yield following applications of biosolids at levels of 7, 18, and 34 dry Mg/ha in the Chihuahuan Desert. Although dormant season applications of biosolids seem to be more beneficial for plant growth than growing season applications during the year of biosolids application (Benton and Wester, 1998), explanations for this phenomenon have not been documented (Mata-Gonzalez *et al.*, 2002).

Most evidence is related to its negative effect on aboveground vegetative and reproductive plant biomass (Hutchings and John, 2003; Milchunas and Lauenroth, 1993), changes in the spatial patterning of plant canopies

and soil resources (Adler *et al.*, 2001; Bertiller and Coronato, 1994; Callaway, 1995; Schlesinger *et al.*, 1990), the reduction of soil seed banks (Bertiller, 1996; Bertiller, 1998), the decrease in the availability of safe micro sites for plant reestablishment (Bisigato, 2000; Oesterheld and Sala, 1990), and the invasion of woody plants (Milchunas and Lauenroth, 1993; Rodriguez *et al.*, 2007; Schlesinger *et al.*, 1990).

Aboveground defoliation can modify the partitioning of assimilates between belowground and aboveground organs and consequently the root growth of defoliated plants (Belsky, 1986; Richards and Caldwell, 1985; Rodriguez *et al.*, 2007; Snyder and Williams, 2003).

In this research we have studied the amount of above ground biomass and occurrence of *Anchusa italica* (*Anchusa azurea*) (Gharaman, 2003) (Fig. 1) at the rangeland area of Shanjan village, Shabestar district, NW Iran. This parameter needs more attention, but it is one of the determining Factors of rangeland ecosystem.

MATERIALS AND METHODS

The research area is part of Shanjan rangeland in Shabestar district with distance about 5 Kilometers from Shabestar city. The terrain in this area is hilly and we carried out the study on a site with a northerly aspect (Bibalani *et al.*, 2011a; Bibalani *et al.*, 2011b) (Fig. 1). This region is component of Iran-Turan Flora with elevation between 1700-1850 m (Bibalani *et al.*, 2011b). This study has been researched in summer 2010.

Anchusa azurea is a species of flowering plant known by the common name Italian bugloss (Fig. 2). It is a



Fig. 1: Part of Shanjan rangeland in Shabestar district, East Azerbaijan province, Iran



Fig. 2.: *Anchusa Italica (Anchusa azurea)* species




Fig. 3: Sampling design in 1*1 m plot (Xiaoyan *et al.*, 2001). 1, 2, 3, 4 and 5 is sub sample in each main sample

bristly perennial which reaches just over half a meter in height. It has straight lance-shaped leaves and petite tubular flowers of bright violet-blue. This species is native to Europe and western Asia but is well-known elsewhere as a noxious weed (Wikipedia, 2011) (Table 1). In this research, Stem biomass has been sampled in May and June, 2010. For sampling, we used an accidental sampling methodology (1*1 m plot) in this research and selected 35 (7 plots with 5 sub sample for each of them) samples in total (Xiaoyan *et al.*, 2001) (Fig. 3).

After sapling from studding area, they have been scaled fresh weight of above ground part of plant with sensitive scale then dried by Avon set in 80°C during

Table 1: Scientific name for *Anchusa italica (Anchusa azurea)* Classification Report (USDA, 2011)

Kingdom	<i>Plantae – Plants</i>
Subkingdom	<i>Tracheobionta – Vascular plants</i>
Superdivision	<i>Spermatophyta – Seed plants</i>
Division	<i>Magnoliophyta – Flowering plants</i>
Class	<i>Magnoliopsida – Dicotyledons</i>
Subclass	<i>Asteridae</i>
Order	<i>Lamiales</i>
Family	<i>Boraginaceae – Borage family</i>
Genus	<i>Anchusa L. – bugloss</i>
Species	<i>Anchusa azurea Mill. – Italian bugloss</i>




Fig. 4: *Anchusa italica (Anchusa azurea)* stem weight (fresh and dried weight)

24 h (Xiaoyan *et al.*, 2001) and scaled dried weight separately. This study have been work in Shanjan rangeland at Shabestar district in East Azerbaijan, Iran in summer 2010.

RESULTS

Results from this study showed that the maximum, minimum and medium stem biomass of *Anchusa italica (Anchusa azurea)* in the study area were 18.5, 31.4 and 22 g, respectively (Fig. 4).

Stem height anchusa italica (*Anchusa azurea*) was unsteady from 350 to 560 mm, that average of it is about 410 mm.

CONCLUSION

In total of 7 plots were identified and 35 samples were studied in this research work. From 35 samples about 84.93% of stem weight was lost when samples were dried.

Vegetation species can have an effect on soil chemical and physical properties (Ardekani, 2003). Increasing *Anchusa italica (Anchusa azurea)* species in the study area could cause specific biological qualification, and as this species increasing density of above ground Biomass will increase, and also the amount of Soil protection and stabling will increase specially

protection with wind erosion and soil lost with runoff (Bibalani *et al.*, 2011a; Bibalani *et al.*, 2011b; Shadkami-Til and Bibalani, 2010; Shadkami-Til and Bibalani, 2011). Study on this plant over ground biomass is so much important information especially for medicinal plant. Joudi and Bibalani (Bibalani *et al.*, 2010) have been studied and recognized some medicinal plant of Ilkhji region, Eastern Azerbaijan Province (Northwestern Iran).

In this study we examined the biomass of this plant and results suggest that changes in the above ground cover of this plant affect by grazing or soil compaction with animal at this area as found in other studies (Bibalani, 2011a; Bibalani, 2011b; Bibalani, 2011c; Bibalani *et al.*, 2010; Bibalani *et al.*, 2011a; Bibalani *et al.*, 2011b; Rodriguez *et al.*, 2007; Shadkami-Til and Bibalani, 2010; Shadkami-Til and Bibalani, 2011) and The difference of wet weight and biomass of this plant would be expected in this area (Bibalani, 2011a; Bibalani, 2011b; Bibalani, 2011c; Bibalani *et al.*, 2010; Bibalani *et al.*, 2011a; Bibalani *et al.*, 2011b; Shadkami-Til and Bibalani, 2011).

This study has revealed and quantified the stem biomass of the *Anchusa italica* (*Anchusa azurea*) in the Shanjan rangelands, the plant has good biomass in this research area and probably also in other areas where the *Anchusa italica* (*Anchusa azurea*) is growing that need studding separately in another areas. It is a pioneer study, and the results have given estimations of the stem biomass of the *Anchusa italica* (*Anchusa azurea*) for the first time in Shanjan rangeland. It is needed for studying this and other shrub species in the area and could be used in identifying plants best suited for rangeland ecosystem stability and specifically for stabilizing surface soil layers especially from water and wind erosion.

ACKNOWLEDGEMENT

The authors greatly acknowledge the scientific support from the Islamic Azad University- Shabestar Branch, to the first author in this study. This paper is a part of a project entitled "Study on Root development forbs and shrubs on Shanjan Range of the Shabestar area, and their effects on soil surface and subsurface erosion" with project number 51955880630001. The authors also express their sincere appreciation to the anonymous reviewer(s) for their help to improve the paper quality.

REFERENCES

Adler, P.B., D.A. Raff and W.K. Lauenroth, 2001. The effect of grazing on the spatial heterogeneity of vegetation. *Oecolog.*, 128: 465-479.
Ardekani, M., 2003. Ecology, University Tehran, Tehran, pp: 340.

Belsky, A.J., 1986. Does herbivory benefit plants? A review of the evidence. *Ameri. Natur.*, 127(6): 870-892.
Benton, M.W. and D.B. Wester, 1998. Biosolids effects on tobosagrass and alkali sacaton in a Chihuahuan desert grassland. *J. Environ. Qualit.*, 27: 199-208.
Bertiller, M.B., 1996. Grazing effects on sustainable semiarid rangelands in Patagonia: The state and dynamics of the soil seed bank. *Environ. Manage.*, 20: 123-132.
Bertiller, M.B. 1998, Spatial patterns of the germinable soil seed bank in northern Patagonia. *Seed Sci. Res.*, 8: 39-45.
Bertiller, M.B. and F. Coronato, 1994. Seed bank patterns of *Festuca pallescens* in semiarid Patagonia (Argentina): A possible limit to bunch reestablishment. *Biolog. Conserv.*, 3: 57-67.
Bibalani, G.H., 2011a. Investigation on persian orange root growth in first tree years. *Int. J. Acad. Res.*, 3(1): 705.
Bibalani, G.H., 2011b. Investigation on prunus avium root growth in first tree years. *Int. J. Acad. Res.*, 3(1): 708.
Bibalani, G.H., 2011c. Investigation on Pyrus Sp. root growth in first tree years. *Int. J. Acad. Res.*, 3(1):122.
Bibalani, G.H., L. Joudi and H. Shadkami-til, 2010. Average stem biomass of lappula microcarpa in Shanjan Rangelands, East Azerbaijan, Iran. *Res. J. Biolog. Sci.*, 5(6): 444.
Bibalani, G.H, L. Joudi and H. Shadkami-til, 2011a. Average stem biomass of paronychia kurdica boiss in Shanjan Rangelands, East Azerbaijan, Iran. *Annal. Biol. Res.*, 2(1): 231.
Bibalani, G.H, L. Joudi and H. Shadkami-til, 2011b. Research of root biomass of lappula microcarpa in Shanjan Rangelands, East Azerbaijan, Iran. *Annal. Biol. Res.*, 2(1): 237.
Bidlock, E.J., J.E. Voughan and C.L. Devald, 1999. Forage quality of 10 estern gama grass. *J. Range Manage.*, 52: 661.
Bisigato, A.J., 2000. Dinamica de la vegetacion en a reas pastoreadas del extremo austral de la provincial fitogeografica del Monte Ph.D. pp: 163.
Callaway, R.M., 1995. Positive interactions among plants. *Botanical. Rev.*, 61: 306-349.
Fresquez, P.R., R.E. Francis and G.L. Dennis, 1990. Soil and vegetation responses to sewage sludge on a degraded semiarid broom snakeweed/blue grama plant community. *J. Range Manage.*, 43: 325-331.
Gharaman, A., 2003. Folor Colored Iran, Froest and Rangland reserch Organization, Tehran.
Hutchings, M.J. and E.A. John, 2003. Distribution of Roots in Soil, and Root Foraging Activity. In: De-Kroon, H. and E.J.W. Visser (Eds.), *Ecological Studies Ecological Studies*, pp: 33-60.

- Mata-Gonzalez, R., E.S. Ronald and W. Changgui, 2002. Shoot and root biomass of desert grasses as affected by biosolids application. *J. Arid Environ.*, 50: 477-488.
- Milchunas, D.G. and W.K. Lauenroth, 1993. Quantitative effects of grazing on vegetation and soils over a global range of environments. *Ecolog. Monogra.*, 63(4): 327-366.
- Mogaaddam, M.R., 2001. Ecology Descriptive and Astistic Vegetal Coverage, Tehran University, Tehran, pp: 258.
- Mozaffarian, V., 2007. A Dicionary of Iranian, Latin, English, Persian, Farhang Moaser, Tehran.
- Oosterheld, M. and O.E. Sala, 1990. Effects of grazing on seedling establishment: The role of seed and safe-site availability. *J. Vegetat. Sci.*, 1: 353-358.
- Richards, J.H. and M.M. Caldwell, 1985. Soluble carbohydrates, concurrent photosynthesis and efficiency in regrowth following defoliation: A field study with *Agropyron* species. *J. Appl. Ecol.*, 22: 907-920.
- Rodriguez, M.V., M.B. Bertiller and C.L. Sain, 2007. Spatial patterns and chemical characteristics of root biomass in ecosystems of the Patagonian Monte disturbed by grazing. *J. Arid Environ.*, 70: 137-151.
- Schlesinger, W.H., J.F. Reynolds, G.L. Cunningham, L.F. Huennke, W.M. Jarrel, R.A. Virgini and W.G. Withford, 1990. Biological feedback in global. *Desertification Sci.*, 247: 1043-1048.
- Shadkami-Til, H. and G.H. Bibalani. 2010. Under-over ground Biomass characterics of perennial Species (*Teucruim polium*) in northwest Iran (Till area of Shabestar) *Int. J. Acad. Res.*, 2(6): 110.
- Shadkami-Til, H. and G.H. Bibalani, 2011. Over ground Biomass characterics of Genera single Species Iran (*Cnicus benedictus*) in northwest Iran (Till area of shabestar). *Int. J. Acad. Res.*, 3(1): 698.
- Snyder, K.A. and D.G. Williams, 2003. Defoliation alters water uptake by deep and shallow roots of *Prosopis velutina* (*Velvet mesquite*). *Funct. Ecolog.* 17:363374.
- USDA., 2011. *Anchusa azurea*. Retrieved from: <http://plants.usda.gov/java/profile?symbol=ANAZ>, (Accessed on: June, 17, 2011).
- Wikipedia, 2011. *Anchusa azurea*. Retrieved from http://en.wikipedia.org/wiki/Anchusa_italica, (Accessed on: June, 17, 2011).
- Xiaoyan, P.G. Zhou, Q. Zhuang, Y. Wang, W. Zuo, G. Shi, X. Lin and Y. Wang, 2001. Effects of sample size and position from monolith and core methods on the estimation of total root biomass in a temperate grassland ecosystem in Inner. Mongol. *Geod.*, 155: 262-268.