

## Research Article

### The Major Factors Affecting Ectomycorrhizal Fungi Diversity in the Forest Ecosystem

Di-Di Chai, Su-Juan Guo, Xiao-Bing Sun and Tian-Tian Qin

Key Laboratory for Silviculture and Conservation in Beijing Forestry University, Beijing

**Abstract:** Ectomycorrhizal fungi are important for plants growth which can enhance their resistance to drought, diseases, infertility and so on in the forest ecosystem, so researches on Ectomycorrhizal fungi are of necessity and great significance. As the basis for Ectomycorrhizal fungi research, the Ectomycorrhizal fungi diversity is attributed to many factors which can be divided to two parts in this review: biotic factors and abiotic factors. Biotic factors include chronosequences, bacteria living with ECM, habitat area, host genotype and abiotic factors include forest fire, toxic metal contamination, nitrogen content, survey method and microenvironment. The effects by these factors are deeply analyzed and discussed based on the plenty of literatures, our previous work and current research. The review not only introduces the current research result, but also arouses new ideas on Ectomycorrhizal fungi diversity research.

**Keywords:** Abiotic factor, biotic factor, diversity, ectomycorrhizal, review

#### INTRODUCTION

Ectomycorrhiza (ECM) which is formed by ectomycorrhizal fungi and the dominant tree roots in the forest ecosystems is widely observed. The ectomycorrhiza improves host performance by enhancing nutrient and water uptake from the soil and protecting host roots from being damaged by salinity, pathogens and toxic compounds (Smith and Read, 1997). Ectomycorrhizal fungi plays an important role in keeping the forest community dynamic in forest ecosystems, so doing research on Ectomycorrhizal fungi is necessary and significant.

The Ectomycorrhizal fungi diversity is the basis for Ectomycorrhizal fungi research, so many studies pay attention to the diversity of Ectomycorrhizal fungi (Dahlberg *et al.*, 2001; Poole *et al.*, 2001; Lilleskov *et al.*, 2001; Koide *et al.*, 2011). Their diversities are complexity, for example, different researches show different diversity though in the same forest composition (Palmer *et al.*, 2008; Ma *et al.*, 2010; Buscardo *et al.*, 2011) or in the same experiment place (Twieg *et al.*, 2009; Richard *et al.*, 2004, 2005), that is owing to too many factors which have an effect on Ectomycorrhizal fungi diversity.

These factors include biotic factors and biotic factors, while most people only focus on a few of them and they do not have a system summary. Every factor may cause the change of Ectomycorrhizal fungi diversity in nature or in the experiment, so reviewing the major factors affecting on Ectomycorrhizal fungi diversity is necessary and significant. This review has

not only introduced the current research result, but also supplied the new ideas on Ectomycorrhizal fungi diversity research.

#### MATERIALS AND METHODS

The method for the aboveground survey of fruiting bodies and the belowground survey of root tipssurveying is vital for identification of ECM fungi, moreover sample time and sample volumes are also important factors. These effects were found in our research, during research on ECM diversity of *Castanea mollissima* in north China, we only found three ECM of *C. mollissima* by the morphological identification in 2010 (Chai *et al.*, 2010), however, by using the molecular identification method and increasing the sample time and sample volumes of fruitbodies collecting, a great more number of ECM species were found living with *C. mollissima* until 2012, so we conclude that survey method is a major factor which affects the ECM diversity of *C. mollissima*.

The aboveground survey of fruiting bodies, morphological classification is vital for surveying aboveground fruiting bodies. If it is used alone, however, it is less effective in identifying community diversity. A combination of morphological classification and molecular analysis is preferable for studying ECM diversity (Dahlberg *et al.*, 2001; Taylor, 2002) and also with continued sequencing of fruiting bodies from aboveground surveys, there should eventually be a reduction in the number of unidentified

**Corresponding Author:** S.J. Guo, Key Laboratory for Silviculture and Conservation, Ministry of Education, Beijing Forestry University, Beijing 100083, China, Tel.: +8613641050478

This work is licensed under a Creative Commons Attribution 4.0 International License (URL: <http://creativecommons.org/licenses/by/4.0/>).

root tip sequences in ECM diversity studies (Horton and Bruns, 2001; Dahlberg *et al.*, 2001), assuming the fruiting body sequences are made available in DNA databases.

The aboveground survey of fruiting bodies is vital for identification of ECM fungi, when combining with the belowground survey; it is a good indicator of community diversity (Gardes and Bruns, 1996; Dahlberg *et al.*, 2001; Horton and Bruns, 2001; Taylor, 2002). The belowground survey of root tips, morphological classification is an effective identification method for belowground surveys (Wurzburger *et al.*, 2001; Sakakibara *et al.*, 2002). However, the results obtained with this method can be affected by poor conservation, phenotypic expression, tip age, environmental conditions, different experimental times and the operators' procedures (Wurzburger *et al.*, 2001; Horton and Bruns, 2001; Sakakibara *et al.*, 2002; Burke *et al.*, 2005). For these reasons, molecular analysis is an ideal identification method for belowground surveys.

In many studies, the aboveground results are not match well with the belowground result. Discrepancies have been explained by the fact that some species rarely or never produce fruiting bodies (Horton and Bruns, 2001), some species form inconspicuous fruiting bodies and therefore are under sampled (Gardes and Bruns, 1996; Tedersoo *et al.*, 2003) and some species (such as the Cantharellales) do not sequence well (Feibelman *et al.*, 1994).

Sample time and sample volumes are also important influences for identification diversity of ECM fungi. In certain ECM diversity studies, a large number of belowground root tip samples were collected only at a single time point. This approach could not reveal the composition of the entire community (Horton and Bruns, 2001; Taylor, 2002). A methodology based on sampling at multiple time points throughout the growing season should be used in the study. As molecular analysis is affected by DNA degradation in storage, processing samples promptly is very important. Small sample volumes should be collected in the study to ensure that the samples could be processed within a week. Palmer used a similar method to identify ECM forming during one growing season (Palmer *et al.*, 2008).

## RESULTS AND DISCUSSION

**Abiotic factors:** ECM root species richness and community structure can be strongly influenced by abiotic factors, including nitrogen content, forest fire, toxic metal contamination, survey method and microenvironment.

**Nitrogen content:** Nitrogen (N) content of soil is the main limiting factor for plants and it can also influence

the diversity of EMF communities (Agerer and Göttslein, 2003), particularly under a deficiency or oversupply of nitrogen (N) (Alexander and Fairley, 1983; Nilsson, 2004; Parrent *et al.*, 2006; Parrent and Vilgalys, 2007). Peter reported that after 2 years of further N supply significant changes were observed. The abundance of species that form large sporocarps (cap diameter about 5-10 cm) decreased on the fine roots in the underground community while the abundance of species with no or resupinate sporocarps increased (Peter *et al.*, 2001a). Similarly, through an increasing anthropogenic N deposition gradient in long-term experiment, species richness of both sporocarp surveys and sampling of the fine roots were decreased (Lilleskov *et al.*, 2001).

However, responses of the relatively rapid decline of sporocarp abundance and diversity to surplus N are different; the belowground community is faster (Lilleskov *et al.*, 2001, 2002a). The diversity of EMF communities in Minnesota (US) was measured by Avis in a 16 year N-addition (fertilization) field experiment (Avis *et al.*, 2003). The evenness and diversity of sporocarp species decreased in the fertilization treatments and total sporocarp species richness was reduced by more than 50%. Carfrae also demonstrated that after a 3-year nitrogen, sulphur and acidity treatment, N deposition suppressed the appearance of EM fungi, producing larger sporocarps in young plantations (Carfrae *et al.*, 2006). Fewer sporocarps and lower EMF sporocarp diversity were generally found under the N treatment plots, while the number of mycorrhizal root tips was greatest. In contrast, Kårén and Wiklund found the opposite; they reported a 50% reduction in sporocarp number and species richness of mycorrhizal species during the N fertilization experiment, while the EMF species richness and diversity on the roots were not changed (Kårén and Nylund, 1997; Wiklund *et al.*, 1995).

Especially, nitrogen should be available and well balanced (Walker *et al.*, 1995; Rygielwicz *et al.*, 1997; Treseder, 2004), but species seem to react differently to supply (Alberton and Kuyper, 2009). Avis found that different species responded differently to the treatment and so the composition of the above-ground ECM community differed across the fertilization treatments. Moreover, they found that the treatment to different species responded differently and the composition of the aboveground EMF community differed. The same response was found in the below-ground result; however, it was smaller than the aboveground result (Avis *et al.*, 2003). N fertilization with ammonium nitrate promoted the growth of *Tomentellopsis submollis*, while inhibited the growth of *Piloderma croceum* mycelium (Arnebrant and Söderström, 1992), this species-specific N uptake capacity under different N availability shows two fungal species have different optimum range of N. A shift in ectomycorrhizal

community structure with reference to N availability has been observed (Fransson *et al.*, 2000, Parrent *et al.*, 2006). The same result that ECM taxa can differ in response to N deposition was found by Lilleskov (Lilleskov *et al.*, 2002b). The reaction to nitrogen may be the different sensitivity of both species to the availability of this nutrient (Alberton and Kuyper, 2009).

To summarise the results of the above fertilisation studies, we can conclude that N surplus is a factor which have an effect on the diversities of ECM community and its effect is sooner in the aboveground response than belowground changes, different EMF taxa can differ in response to N deposition. Therefore, a refined approach with nitrogen amendment adjusted to the special demands of either species would be necessary.

**Forest fire:** Forest fire is the major forest disturbance pattern and hazard (Byrot, 2009) which can affect ECM fungal communities (Grogan *et al.*, 2000; Chen and Cairney, 2002; Anderson *et al.*, 2007) by the cessation of carbohydrate flow to the established, native ECM fungal population. However, it is difficult to predict the extent to the forest fire affects the population of ECM fungi as there are multiple factors (Koide *et al.*, 2011). For example, after fire changes in the chemical or biological properties of the soil (Amaranthus *et al.*, 1993; Colinas *et al.*, 1994) and the presence of plant species which can resprout or quickly establish (Borchers and Perry, 1990)? However, some environmental factors are not affected by the fire, belowground soil temperature during the fire probably did not rise beyond 200°C. Consequently, concentration of phosphorous which would be affected by even higher temperatures than nitrogen remained almost unchanged (Knoepp *et al.*, 2005). Soil pH was slightly more alkaline at post-fire sites, a common effect in burnt soils and in this case probably mainly due to the increased concentration of soluble calcium which is a product of ash deposition (Knoepp *et al.*, 2005).

Though the fire influence is complex, we still found some regularity that the frequency of the fire is a major factor for ECM diversity changing.

Many studies on fire influence show that low-frequency fire events do usually not cause substantial change but the relative abundance of ECM. The result of the ECM community of the *Quercus ilex* forest at Nazar between 1998 and 2001 demonstrated that there are differences in the relative abundance of certain ECM morphotypes when comparing the stand affected by fire and the undisturbed plot (Román and Miguel, 2005). Anderson study changes in vertical stratification of soil basidiomycetes by molecular tools, result showed no substantial change in species richness (Anderson *et al.*, 2007). A similar trend was observed by Tuininga on ectomycorrhizal roots of adult pine

trees previously exposed to low-frequency, they found no consistent stand-specific changes in fungal diversity but clear shifts in morphotype richness between soil horizons compared to unburned plots (Tuininga and Dighton, 2004). Compared ectomycorrhizal morphotype richness on seedlings from the burnt sites with the control seedlings, only a slight small richness effect by low-frequency fire (Palfner *et al.*, 2008). Fungal species diversity is often not substantially changed by low-frequency fire, which has been reported generally (Torres and Honrubia, 1997; Anderson *et al.*, 2007; Tuininga and Dighton, 2004). Similar results have been reported in the ECM community of a *Pinus sylvestris* forest in Sweden (Jonsson *et al.*, 1999) and a *Picea* forest in Canada (Mah *et al.*, 2001). A decrease in the number of mycorrhizal tips after low-frequency fire has also been reported by several studies (Dahlberg *et al.*, 2001; Stendell *et al.*, 1999; Torres and Honrubia, 1997). It was concluded that low-frequency fire induced a shift in the relative abundance of each species rather than a change in the species composition.

In contrast, several published studies report that the intensity fire results differ in several aspects from the above results. For example, after the intensity fire in a *Pinus muricata* forest in California, a shift in the species composition of the ECM community were found by Baar and Dahlberg found both a decrease and a change in the species composition in the ECM diversity of several conifer forests affected by fire in Scandinavia (Baar *et al.*, 1999; Dahlberg *et al.*, 2001). The intensity of fire is an important factor which would influence the impact of fire on the ECM community (Dahlberg *et al.*, 2001; Dahlberg, 2002; Grogan *et al.*, 2000).

Many trees survive and the organic matter remains intact after a low-intensity fire and this situation causes small affect of the fire on the ECM community. Loss of fungal species richness under the ground but may alter community structure. However, high burning frequency may lead to a loss of ectomycorrhizal fungal diversity as reported by Bastias (Bastias *et al.*, 2006).

After fire influence, the diversities of ectomycorrhizal fungi are finding in the forest and most of them are belonged to Ascomycota and Basidiomycetes.

Most studies are about Ascomycota fungi. Post-fire fungi which is a series of typically fungi appeared only after a fire, largely belonging to the order Pezizales and these fungi produced fruiting nearly 6 weeks after a fire (Petersen, 1970). Post-fire Pezizales are often referred to as facultatively mycorrhizal. For example, *Biotrophism* which belongs to Pezizalean is supported by direct observation of the fungi colonizing root tips in vitro (Warcup, 1990; Dahlstrom *et al.*, 2000) as well as in vivo (Vrålstad *et al.*, 1998). Some species of post-fire Pezizales are reported to form ectendomycorrhizal

associations with members of the *Pinaceae* (Danielson, 1984; Egger and Paden, 1986; Dahlstrom *et al.*, 2000). *Wilcoxina* sp. is common found on post-fire pine (*Pinus muricata*) seedlings (Baar *et al.*, 1999). *Geopora cooperi* is widely distributed in undisturbed forest conditions (Maia *et al.*, 1996; States and Gaud, 1997) as well as in nutrient-poor, volcanic cinder soils with species of the *Pinaceae* (Gehring *et al.* 1998). *Wilcoxina rehmii* and *Geopora cooperi* belongs to the Pezizalean species, the probability of recovery of them have been increased because the fire removed most of the speciose mycorrhizal fungal community that typically occurs in undisturbed habitats (Horton *et al.*, 1998; Baar *et al.*, 1999; Stendell *et al.*, 1999; Fujimura *et al.*, 2005). They found very low richness of mycorrhizae, mainly represented by a hymenoscyphus-like morphotype and *Cenococcum geophilum*, both of them were more abundant on seedlings from unburnt forest than on those from the burnt areas (Palfner *et al.*, 2008). Ascomycota mycorrhizal may be less abundant on post-fire seedlings than on trees not affected by fire (Izzo *et al.*, 2006; Fujimura *et al.*, 2005). The study on the ecosystem function of *Morchella* showed that some species fruiting may be mycorrhizal without a fire influence (Dahlstrom *et al.*, 2000; Hobbie *et al.*, 2001), however, they may not form mycorrhizas in burned sites, if they do, they may revert to being saprobic (Buscot, 1994; Dahlstrom *et al.*, 2000). Isotopic experiment on *Morchella* species from burned and nonburned sites may help distinguish between mycorrhizal and saprobic functions of post-fire *Morchella* species (Hobbie *et al.*, 2001, 2002). Every year, the wildfires of Mediterranean cause the loss of forested lands where soil and climatic characteristics are suitable for *truffle* cultivation. After the forest fires the competitiveness of *Tuber melanosporum* was highlighted within the ECM community in these soils (Aragón *et al.*, 2012).

Basidiomycetes were the most abundant fungi on both plant groups (post-fire and control), with a clear dominance of a single mycorrhizal morphotype, especially on the post-fire sites with almost double as many (42.7%) colonized root tips as on control seedlings (22.1%) (Palfner *et al.*, 2008). The studies by Palfner and Godoy and Palfner already yielded some evidence that mycorrhizae of *Descolea flavoannulata* Horak are especially abundant on seedlings or young trees at disturbed or ruderal sites: they were found in such distinct habitats as on seedlings of *Nothofagus pumilio*, establishing on thick ash layers in Patagonia in the aftermath of a volcanic eruption, on 2-year old nursery seedlings of *Nothofagus alpina* and on planted trees of the same species in park-like environments (Palfner, 1997; Palfner, 2001; Godoy and Palfner, 1997). The less frequent fungal species such as *Cortinarius magellanicus* whose mycorrhiza is characterized by conspicuous pseudosclerotia, russuloid morphotypes

and tomentelloid morphotypes (Palfner, 2001; Palfner *et al.*, 2002), the first morphotypes were found only on seedlings from the undisturbed control area and the other two morphotypes were appeared almost exclusively on roots from the burnt sites. It should be mentioned that Bastias also report absence of Cortinariaceae in the aboveground horizon of a frequently burnt *Eucalyptus pilularis* forest (Bastias *et al.*, 2006). Increased richness of a single fungal species or species complex on mycorrhizal roots after the fire influence has already been studied: 6 years after the fire event, *Suillus brevipes* were found the most abundant species associated with *Pinus banksiana* (Visser, 1995), early stage after forest fire, *Rhizopogon* spp. was dominant colonizers of bioassay seedlings of *Pinus muricata* and *Pinus jeffreyi*, respectively fruiting patterns of macromycetes often revealed a boost of ascomycetous taxa (Baar *et al.*, 1999; Izzo *et al.*, 2006).

**Heavy metal contamination:** It was shown that the metal-contaminated soils had a negative impact on the abundance and variability of ectomycorrhizas in the root systems.

Different kinds of Heavy Metal (HM) environment showed different diversities of ECM, these effects have been found in the birch root systems (Bojarczuk and Barbara, 2010). Under natural conditions, *Betula pendula* develops Ectomycorrhizal (ECM) symbiosis (Smith and Read, 1997). *Betula pendula* often has been found on soils contaminated with metals, such as Zn (Denny and Wilkins, 1987); Pb (Eltrop *et al.*, 1991); Zn and Pb (Kopponen *et al.*, 2001); Cd, Cd and Zn (Regvar *et al.*, 2006); and Cu and Pb (Klink *et al.*, 2006), however, the diversity of Ectomycorrhizal fungi are not same in each place. The reason is that ECM diversity has different response to the different HM type. However, the same ectomycorrhizal fungi when hosted different trees has also different response to the same HM type. For example, the phytoextraction capacity of willow or poplar associated with mycorrhiza has been studied by Lingua (Lingua *et al.*, 2008). Lingua reported that *Populus alba* (clone Villafranca) inoculated with *Glomus mosseae* or *Glomus intraradices* accumulated lower Zn concentrations; *Populus nigra* (clone Jean Pourtet) also accumulated lower Zn concentrations when inoculated with *G. mosseae* whereas its Zn concentrations remained unchanged when inoculated with *G. intraradices* (Lingua *et al.*, 2008). On the other hand, Sell reported that the Cd concentration were enhanced in *Populus canadensis* when associated with an ectomycorrhizal fungal species (*Paxillus involutus*), however, when associated with this ectomycorrhizal fungi, the Cd concentration remained unchanged in *Salix viminalis* (clone 78198) (Sell *et al.*, 2005). By contrast, Baum reported higher Cd, Zn and Cu concentrations in the

stems of *Salix dasyclados* (clone SDHM) associated with the same ectomycorrhizal fungi (Baum *et al.*, 2006).

Living at high metal concentrations causes a high variability in tolerance to ECM fungal species. (Fomina *et al.*, 2005; Leski *et al.*, 1995). The effects of heavy metals are inconsistent to different ECM as a result of this fungal variability (Godbold *et al.*, 1998; Meharg and Cairney, 2000). Metal enriched soils can lead to the selection pressure of metal toxicity ECM genotypes (Hartley *et al.*, 1997; Leyval *et al.*, 1997; Markkola *et al.*, 2002; Colpaert *et al.*, 2004; Adriaensen *et al.*, 2005). ECM has the ability to alleviate the effects of heavy metal toxicity for themselves and their host trees (Adriaensen *et al.*, 2005, 2006; Krznanic *et al.*, 2009), by providing a more balanced access to mineral nutrition (Marschner and Dell, 1994).

The kinds of ECM which are always found in HM polluted soils are mainly Ascomycete. For example, on the roots of *Salix caprea* studied at the polluted site 15 ECM taxa were found by combining the ECM phenotyping and sporocarp identification, the members of Sordariaceae were the most frequent, followed by The lephoraceae (Regvar, 2010). A study from a primary succession site at the forefront of a receding glacier also reported Sordariaceae as dominant ECM fungi of willows (Trowbridge and Jumpponen, 2004). *Glomus mosseae* which has frequently been identified in HM soil (Hassan *et al.*, 2009; Vallino *et al.*, 2006), these ascomycetes could be intercellular endophytes or rhizoplane colonisers and their role for hosts deserve further attention. Ascomycete ECM fungal taxa seem to be most numerous at the plots with high metal and low levels of organic matter and more sparse vegetation cover.

**Survey method:** This part was in material sand methods.

**Microenvironment:** ECM living environment which has great effect on the diversity of ECM are complex and hard to control, the research on microenvironment factors can help us clearly know the ECM living habit. Vitro environment is the best way to realize microenvironment, our laboratory is focus on the vitro of *C. mollissima*, we have already built up the regeneration system of *C. mollissima* (Hou *et al.*, 2010), three new ECM species were separating from the *C. mollissima* ectomycorrhiza and two of them were successful infecting with the roots of *C. mollissima* in vitro environment. During the whole process of growth, we found the vitro environment factors have different influence on different kind of *C. mollissima* ECM which include carbon dioxide concentration, relative humidity, PH value, growth hormone levels and so on, these factors which affect the ECM diversity of *C. mollissima* are our new current research.

**Biotic factors:** ECM root species richness and community structure are also influenced by a range of biotic factors, including chronosequences; bacteria lived with ECM, host genotype, habitat area and so on.

**Chrono sequences:** The age of plants living with ECM is an important factor to show their nutritional status and the structure, which are correlated with the ECM diversity. So the chronosequence affects the ECM species compositions. For instance, Peter showed different EMF species compositions were found in different stand ages (35, 100–200 years old) (Peter *et al.*, 2001b). They explained that the sites with various ages may lead to their results. The same study was carried out by Visser, in which he indicated that sequence of mycorrhizal fungi related to stand age has undergone regeneration following wildfire disturbance in both fruiting body and root tip survey (Visser, 1995). Visser found different diversities in different stand ages, so he divided mycorrhizal fungi into early-stage, middle-stage and late-stage fungi (Visser, 1995).

Different experiment showed different results, most people believe that as the ages increase ECM diversities increase too, however some people showed that as the ages increase the diversities of EMF decrease, there are still some researchers found that ECM diversity are different though in the same ages.

As the ages increase ECM diversities increase. Our laboratory is major research on the ECM diversity of chestnut (*Castanea mollissima*) in north China, Chronosequence factor which affects the ECM diversity of *C. mollissima* are our current research. During the long term research, our sample plot included 15 years old *C. mollissima* stand and over 80 years *C. mollissima* stand, both aboveground and belowground diversities were much more found in the over 80 years stand, so we believe that as the *C. mollissima* forest ages increase their ECM diversities increase too. A progressive increase in species richness of mycobionts along chronosequences of Sitka spruce forests (6, 12, 30, 40 years) in the result of both sporocarp survey and root tip sampling has been demonstrated by Palfner (Palfner *et al.*, 2005). Gebhardt showed that the diversity of ECM sporocarps and root tip morphotypes of red oak growing on forest reclamation sites (Gebhardt *et al.*, 2007). Each stand (5, 21, 33, 43 and 46 years old) results were demonstrated stand specific EMF communities with low similarity to other stands by both methods. The total number of EM species was the largest in the 46 year-old nature stand in both sporocarp survey and the root sampling. ECM fungal community structure was surveyed in four combinations of age class (5-, 26-, 65- and 100-year-old), each representing critical seral stages of stand development in Interior Cedar-Hemlock (ICH) forests of southern British Columbia. ECM fungal community structure was more strongly influenced by stand age and in particular, no

pattern that paralleled the strong increase in ECM fungal diversity seen from 5-year-old to 26-year-old age classes (Twieg *et al.*, 2009). Tabea Kipfer investigated the diversity of ECM fungal communities on a chronosequence of 12 *Pinus sylvestris* stands in Valais (Switzerland) and Val d'Aosta (Italy) between 1990 and 2006, the number of ECM species was significantly lower in samples from recently (2-5 years) (Kipfer *et al.*, 2011).

As the ages increase, the diversities of EMF decrease. Species richness decreases as the age of host increases in the results of fruiting body surveys, however, the same result were not found in the results of belowground sampling (Richard *et al.*, 2004, 2005). The high diversity of mycorrhizal fungal taxa existing in seedlings also confirms the applicability of using seedlings to effectively document mycobiont diversity in situ (Walker *et al.*, 2008). Ectomycorrhizal (ECM) fungal communities of *Quercus liaotungensis* of different ages (seedlings, young trees and mature trees) in the growing seasons between 2007 and 2009 were studied by Wang in a temperate forest of northern China. He indicated that oak seedlings in natural forests are associated with a wide assemblage of ECM fungi as they are with the mature oak trees (Wang *et al.*, 2012).

ECM diversities are different though in the same stand age. Differences in mycorrhizal fungus communities on roots of same-aged seedlings across a deglaciated chronosequence was evaluated by Helm, overall, the EM formation on transplants during the first 2 years was lower and less diverse than on naturally occurring plants in this same chronosequence (Helm and Allen, 1999). Moreover, chronosequence affects are not limited by composition of forest, Twieg described the species composition of the ECM fungal community changed along a chronosequence of mixed forest stands that were similar in vegetation composition and site quality (Twieg *et al.*, 2007).

**Ectomycorrhiza Associated Bacteria (EMAB):** The establishment of plant-fungi interactions may be influenced by bacteria that occur on the surface of mycorrhizal roots and at intercellular and intracellular locations within the ectomycorrhizal mantle and Hartig net (Nurmiaho-Lassila *et al.*, 1997; Mogge *et al.*, 2000). Parts of the bacteria which positively affect the formation of ectomycorrhizal associations were called EMAB (Garbaye, 1994) and these associated bacteria can enhance the symbiosis between host plants and mycorrhizal fungi and support mycorrhiza formation or function (Dunstan *et al.*, 1998; Gryndler *et al.*, 2000; Duponnois and Plenchette, 2003; Kataoka and Futai, 2009). So far, positive effect of EMAB on the mycorrhiza formation and indirect effect of mycorrhiza diversity have been described in many publications (Garbaye, 1994; Poole *et al.*, 2001; Hryniewicz *et al.*, 2009).

Different EMAB have different influence on stimulating mycorrhization and mycorrhiza diversity. For example, mycorrhization of *Amanita rubescens* or *Hebeloma sinapizans* by dual inoculating with EMAB *Pseudomonas putida* or *Bacillus cereus* on *Pinus sylvestris* seedling growth, the result showed that the increased growth of pine seedlings was especially seen for co-inoculation with *P. putida*, *Amanita rubescens* was more efficient in this stimulation than *H. sinapizans*, as *H. sinapizans* was hard to form ECM without the help of its associated EMAB and this may cause changes in mycorrhiza diversity (Kozdrój *et al.*, 2007). The same report has been found that *A. rubescens* mycorrhization was hard to format in the presence of EMAB bacterial strains inoculation by Shishido (Shishido *et al.*, 1996). The result indicates that bacteria only affect mycorrhization when they are associated with mycorrhiza, therefore mycorrhiza diversity is affected by their associated bacteria. Pine seedlings co-inoculation with some ECM was greatly stimulated by *P. putida* of EMAB, many pseudomonads and bacilli promoting kinds of ECM formation have been identified as bacteria (Frey-Klett *et al.*, 1997; Poole *et al.*, 2001).

Explanation of functional interactions between ectomycorrhizal symbionts and their EMAB is still a big challenge. Probanza reported that some strains of *Bacillus*, belonging to EMAB, directly affected the growth of mycorrhiza by gibberellin production (Probanza *et al.*, 2002). de Boer found a possible mechanism for selection of fungus associated bacterial strains by ectomycorrhizal fungi which could select antibiotic-resistant bacteria by exudation of soluble fungal storage sugars, polyols or organic acids which can increase the number of bacteria or exudation of inhibitory chemicals (wiestse *et al.*, 2005).

**Host genotype:** ECM species richness and diversity are correlated with host plant diversity, the effect on vegetation composition forest ecosystem is not the same with that on mixed forest ecosystem.

In vegetation composition forest ecosystem, different hosts have different communities of ECM, taking broad leaved tree species and conifer for example, Palmer investigated ectomycorrhizal community of an American chestnut, finding that only ten putative ECM species were definitively associated with *Castanea dentate* (Palmer *et al.*, 2008), however, Ma found that 22 ectomycorrhizal fungal species comprised the ECM community of *Pinus dens flora* (Ma *et al.*, 2010). The differences were also found inside of broad leaved tree species and conifer, Palmer found that the ECM composition and diversity of *Quercus* spp. are different from that of *Castanea dentate* (Palmer *et al.*, 2008). A total of 30 taxa colonized with *Pinus pinaster* were found by Buscardo, which is more than that with *Pinus densiflora* (Buscardo *et al.*, 2011).

In mixed forest ecosystem, sporocarp inventory result showed that fruiting of some fungal species near either *Q. ilex* or *Arbutus unedo* and this phenomenon was also found in the belowground survey (Richard *et al.*, 2004). EMF species richness and diversity is greatly different from the results in vegetation composition forest ecosystem, so in a mixed forest ecosystem, identification of host plants is imperative for studying EMF species richness and diversity (Richard *et al.* 2004).

**Habitat area:** Several studies have reported that habitat area affects the richness and diversity of ECM fungi, this affect is complex and interesting.

Some people believe that as the habitat area increase the richness and diversity of ECM fungi are also increase, for instance, Peay found that island size had a strong effect on diversity of ECM (Peay *et al.*, 2007). Total species richness increased significantly with island area, independently of the approach used. *Pinus densiflora* Sieb. is one of the major ectomycorrhizal host trees in the eastern Asian biome. Twenty one species of ECM fungi fruit bodies were reported from a 700 m<sup>2</sup> study area (Fujita *et al.*, 1989) while 63 species were found in a 3200 m<sup>2</sup> study area in *P. densiflora* forests in one year based on sporocarp production (Yamada and Katsuya, 1995). Four different study areas found different diversity within a 7,000 km<sup>2</sup> area New South Wales (Hitchcock *et al.*, 2011).

On the opposite, some people found that as the habitat area increase the richness and diversity of ECM fungi are decrease. For example, Durall (Durall *et al.*, 1999) studied the effects of small forest gaps and partial cutting on diversity of EM mushroom, results showed that sporocarp species richness were decreased as gap area increased along 100 m long transects. ECM richness on seedling root tips also decreased slightly with increasing distance from the edge of the intact forest. The maximum richness was found to be 7 m or less from the forest edge for both tree species investigated.

## CONCLUSION

In this study, the abiotic and biotic factors effect on the ECM root species richness and community were discussed. Among the abiotic factors, the effect of N surplus is sooner in the aboveground response than belowground changes; low-frequency fire events do usually not cause substantial change but the relative abundance of ECM. Vitro environment factors, different carbon dioxide concentration, relative humidity, PH value and different growth hormone levels have different influence on the diversity of ECM.

The biotic factors effect on the ECM root species richness and community were also discussed. In many cases, ECM diversities usually increase with hosts aging. Effect of EMAB on the mycorrhiza formation and indirect effect of ECM diversity are positive;

identification of host plants is imperative for studying EMF species richness, especially in mixed forests; the method for aboveground survey of fruiting bodies and the belowground survey of root tips is vital for identification of ECM fungi, moreover sample time and sample volumes are also important.

The knowledge of ECM diversity are more comprehensive by learning more affecting factors, so further studies on the diversity of ECM are still needed in the future.

## ACKNOWLEDGMENT

This study was sponsored by National forestry bureau public industry research and development of special projects “Chestnut industry chain environment friendly production key technology research and demonstration” (201204401); “Twelfth Five-Year” technology support subject “Key technology research and demonstration in high-efficiency production of Chestnut”(2013BAD14B0402).

## REFERENCES

- Adriaensen, K., T. Vralstad, J.P. Noben, J. Vangronsveld and J.V. Colpaert, 2005. Copper-adapted *Suillus luteus*, a symbiotic solution for pines colonizing cu mine spoils. Appl. Environ. Microbiol., 71(11): 7279-7284.
- Adriaensen, K., J. Vangronsveld and J.V. Colpaert, 2006. Zinc-tolerant *Suillus bovinus* improves growth of Zn-exposed *Pinus sylvestris* seedlings. Mycorrhiza, 16(87): 553-558.
- Agerer, R. and A. Göttlein, 2003. Correlations between projection area of ectomycorrhizae and H<sub>2</sub>O extractable nutrients in organic soil layers. Mycol. Prog., 2: 45-52.
- Alberton, O. and T.W. Kuyper, 2009. Ectomycorrhizal fungi associated with *Pinus Sylvester's* seedlings respond differently to increased carbon and nitrogen availability: Implications for ecosystem responses to global change. Glob. Chang Biol., 15(1): 166-175.
- Alexander, I.J. and R.J. Fairley, 1983. Effects of N fertilization on populations of fine roots and mycorrhizas in spruce humus. Plant Soil, 71: 49-54.
- Amaranthus, M.P., J.M. Trappe and D.A. Perry, 1993. Soil moisture, native revegetation and pinus lambertiana seedling survival, growth and mycorrhiza formation following wildfire and grass seeding. Restore Ecol., 1: 188-195.
- Anderson, I.C., B.A. Bastias, D.R. Genney, P.I. Parkin and J.W.G. Cairney, 2007. Basidiomycete fungal communities in Australian sclerophyll forest soil are altered by repeated prescribed burning. Mycol. Res., 111: 482-486.

- Arnebrant, K. and B. Söderström, 1992. Effects of fertilizer treatments on ectomycorrhizal colonization potential in two Scots pine forests in Sweden. *Forest Ecol. Manag.*, 53(1-4): 77-89.
- Aragón, J.M., C. Fischer, J.A. Bonet, A. Olivera, D. Oliach and C. Colinas, 2012. Economically profitable post fire restoration with blacktruffle (*Tuber melanosporum*) producing plantations. *New Forests*, 43: 615-630.
- Avis, P.G., D.J. McLaughlin, B.C. Dentinger and P.B. Reich, 2003. Long term increase in nitrogen supply alters above- and below-ground ectomycorrhizal communities and increases the dominance of *Russula* spp. in temperate oak savanna. *New Phytologist*, 160: 239-253.
- Baar, J., T.R. Horton, A.M. Kretzer and T.D. Bruns, 1999. Mycorrhizal colonization of *Pinus muricata* from resistant propagules after a stand-replacing wildfire. *New Phytologist*, 143: 409-418.
- Bastias, B.A., Z. Xu and W.G. Cairney, 2006. Influence of long-term repeated prescribed burning on mycelial communities of ectomycorrhizal fungi. *New Phytologist*, 172(1): 149-158.
- Baum, C., K. Hryniewicz, P. Leinweber and R. Meißner, 2006. Heavy metal mobilization and uptake by mycorrhizal and nonmycorrhizal willows. *J. Plant Nutr. Soil Sci.*, 169: 516-522.
- Bojarczuk, K. and K.R. Barbara, 2010. Effect of ectomycorrhiza on Cu and Pb accumulation in leaves and roots of silver birch (*Betula pendula* Roth.) seedlings grown in metal-contaminated soil. *Water Air Soil Pollut.*, 207(1-4): 227-240.
- Borchers, S.L. and D.A. Perry, 1990. Growth and ectomycorrhiza formation of Douglas-fir seedlings grown in soils collected at different distances from pioneering hardwoods in southwest Oregon clear-cuts. *Can. J. Forest Res.*, 20(6): 712-721.
- Buscot, F., 1994. Ectomycorrhizal types and end bacteria associated with ectomycorrhizas of *Morchella elata* (Fr.) Boudoir with *Picea abies* (L.) Karst. *Mycorrhiza*, 4(5): 223-232.
- Burke, D.J., K.J. Martin, P.T. Rygiel and M.A. Topa, 2005. Ectomycorrhizal fungi identification in single and pooled root samples: Terminal Restriction Fragment Length Polymorphism (TRFLP) and morph typing compared. *Soil Biol. Biochem.*, 37: 1683-1694.
- Buscardo, E., H. Freitas, J.S. Pereira and P.D. Angelis, 2011. Common environmental factors explain both ectomycorrhizal species diversity and pine regeneration variability in a post-fire Mediterranean forest. *Mycorrhiza*, 21(6): 549-558.
- Byrot, Y., 2009. Living with wildfires: What science can tell us? *Eur. Forest Inst.*, 15: 82.
- Carfrae, J.A., K.R. Skene, L.J. Sheppard, K. Ingleby and A. Crossley, 2006. Effects of nitrogen with and without acidified sulphur on an ectomycorrhizal community in Sitka spruce (*Picea sitchensis* Bong. Carr) forest. *Environ. Pollut.*, 141: 131-138.
- Hitchcock, C.J., S.M. Chambers and W.G. John, 2011. Genetic population structure of the ectomycorrhizal fungus *Pisolithus microcarpus* suggests high gene flow in south-eastern Australia. *Mycorrhiza*, 21(2): 131-137.
- Chen, D.M. and J.W.G. Cairney, 2002. Investigation of the influence of prescribed burning on ITS profiles of ectomycorrhizal and other soil fungi at three Australian sclerophyll forest sites. *Mycol. Res.*, 106: 532-540.
- Chai, D.D., S.J. Guo and X.D. Niu, 2010. Mycorrhizal fungi of wild chestnut isolation and the effect of backing up. *North. Hortic.*, 21: 18-21.
- Colinas, C., D. Perry, R. Molina and M. Amaranthus, 1994. Survival and growth of *Pseudotsuga menziesii* seedlings inoculated with biocide-treated soils at planting in a degraded clear-cut. *Can. J. Forest Res.*, 24: 1741-1749.
- Colpaert, J.V., L.A.H. Muller, M. Lambaerts, K. Adriaensen and J. Vangronsveld, 2004. Evolutionary adaptation to Zn toxicity in populations of Suilloid fungi. *New Phytol.*, 162: 549-559.
- Dahlberg, A., J. Schimmel, A.F.S. Taylor and H. Johannesson, 2001. Post-fire legacy of ectomycorrhizal fungal communities in the Swedish boreal forest in relation to fire severity and logging intensity. *Biol. Conserv.*, 100: 151-161.
- Dahlberg, A., 2002. Effects of fire on ectomycorrhizal fungi in Fennoscandian boreal forests. *Silva Fenn.*, 36(1): 69-80.
- Dahlstrom, J.L., J.E. Smith and N.S. Weber, 2000. Mycorrhiza-like interaction by *Morchella* with species of the Pinaceae in pure culture. *Mycorrhiza*, 9: 272-279.
- Danielson, R.M., 1984. Ectomycorrhizae formation by the operculate discomycete *Sphaerospora brunnea* ezizales. *Mycologia*, 76: 454-461.
- Denny, H.J. and D.A. Wilkins, 1987. Zinc tolerance in *Betula* species. I. effect of external concentrations of zinc on growth and uptake. *New Phytologist*, 106: 517-524.
- Wietse, D.B., L.B. Folman, R.C. Summerbell and L. Boddy, 2005. Living in a fungal world: Impact of fungi on soil bacterial niche development. *FEMS Microbiol. Rev.*, 29: 795-811.
- Dunstan, W.A., N. Malajczuk and B. Dell, 1998. Effects of bacteria on mycorrhizal development and growth of container grown *Eucalyptus diversicolor* F. Mull seedlings. *Plant Soil*, 201(2): 241-249.
- Durall, D.M., M.D. Jones, E.F. Wright, P. Kroeger and K.D. Coates, 1999. Species richness of ectomycorrhizal fungi in cut blocs of different size in the interior cedar-hemlock forest of northwestern British Columbia: Sporocarp and ectomycorrhizae. *Can. J. Forest Res.*, 29: 1322-1332.



- Egger, K.N. and J.W. Paden, 1986. Biotrophic associations between *lodgepole pine* seedlings and post-fire ascomycetes (Pezizales) in monoxenic culture. *Can. J. Bot.*, 64: 2719-2725.
- Eltrop, L., G. Brown, O. Joachim and K. Brinkmann, 1991. Lead tolerance of betula and salix in the mining area of Metternich/Germany. *Plant Soil*, 131: 275-285.
- Feibelman, T., P. Bayman and G. William, 1994. Length variation in the internal transcribed spacer of ribosomal DNA in chanterelles. *Mycol. Res.*, 98(6): 614-618.
- Fomina, M., K. Ritz and G.M. Gadd, 2005. Negative fungal chemotropism to toxic metals. *FEMS Microbiol. Lett.*, 193: 207-211.
- Frey-Klett, P., J.C. Pierrat and L. Garbaye, 1997. Localization and survival of mycorrhizal helper *Pseudomonas fluorescens* during establishment of ectomycorrhizal symbiosis between *Laccaria bicolor* and douglas-fir. *Appl. Environ. Microbiol.*, 63: 139-144.
- Fransson, P.M.A., A.F.S. Taylor and R.D. Finlay, 2000. Effects of continuous optimal fertilization on belowground ectomycorrhizal community structure in a Norway spruce forest. *Tree Physiol.*, 20: 599-606.
- Fujimura, K.E., J.E. Smith, T.R. Horton, N.S. Weber and J.W. Spatafora, 2005. Pezizalean mycorrhizas and sporocarps in ponderosa pine (*Pinus ponderosa*) after prescribed fires in eastern Oregon, USA. *Mycorrhiza*, 15: 79-86.
- Garbaye, J., 1994. Helper bacteria: A new dimension to the mycorrhizal symbiosis. *New Phytol.*, 128: 197-210.
- Gardes, M. and T.D. Bruns, 1996. Community structure of ectomycorrhizal fungi in a *Pinus muricata* forest: Above- and below-ground views. *Can. J. Bot.*, 74: 1572-1583.
- Gebhardt, S., K. Neubert, B. Münzenberger and R.F. Hüttel, 2007. Ectomycorrhiza communities of red oak (*Quercus rubra* L.) of different age in the lusatia lignite mining district. *East Germany Mycorrhiza*, 17: 279-290.
- Gehring, C.A., T.C. Theimer, T.G. Whitham and P. Keim, 1998. Ectomycorrhizal fungal community structure of pinyon pines growing in two environmental extremes. *Ecology*, 79: 1562-1572.
- Godbold, D.L., G. Jentschke, S. Winter and P. Marschner, 1998. Ectomycorrhizas and amelioration of metal stress in forest trees. *Chemosphere*, 36: 757-762.
- Grogan, P., J. Baar and T.D. Bruns, 2000. Below-ground ectomycorrhizal community structure in a recently burned bishop pine forest. *J. Ecol.*, 88: 1051-1062.
- Hassan, S.E., M. Hijri and M. St-Arnaud, 2009. PCR-DGGE analysis of arbuscular mycorrhizal fungi diversity in heavy metal polluted soils. *Proceedings of the 59th Ann Conference Can Soil Microbial*, Concordia University, Montreal, Canada, pp: 69.
- Hartley, J., J.W.G. Cairney and A.A. Meharg, 1997. Do ectomycorrhizal fungi exhibit adaptive tolerance to potentially toxic metals in the environment? *Plant Soil*, 189: 303-319.
- Helm, D.J. and J.M. Allen, 1999. Plant growth and ectomycorrhiza formation by transplants on deglaciated land near exit glacier, Alaska. *Mycorrhiza*, 8: 297-304.
- Hobbie, E.A., N.S. Weber and J.M. Trappe, 2001. Determining mycorrhizal or saprotrophic status of fungi from isotopic evidence: Implications for element cycling and fungal evolution. *New Phytol.*, 150: 601-610.
- Hobbie, E.A., N.S. Weber, J.M. Trappe and G.J. Klinton, 2002. Using radiocarbon to determine the mycorrhizal status of fungi. *New Phytol.*, 156: 129-136.
- Horton, T.R., E. Cázares and T.D. Bruns, 1998. Ectomycorrhizal, vesicular-arbuscular and dark septate fungal colonization of bishop pine (*Pinus muricata*) seedlings in the first five months of growth after wildfire. *Mycorrhiza*, 8: 11-18.
- Horton, T.S. and T.D. Bruns, 2001. The molecular revolution in ectomycorrhizal ecology: Peeking into the black-box. *Mol. Ecol.*, 10: 1855-1871.
- Hou, J.W., S.J. Guo and G.Y. Wang, 2010. Effects of in vitro subculture on the physiological characteristics of adventitious root formation in micro shoot of Castaneda mollissima cv 'yanshanhong'. *J. Forestry Res.*, 21: 155-160.
- Hryniewicz, K., C. Baum, J. Niedojadło and H. Dahm, 2009. Promotion of mycorrhiza formation and growth of willows by the bacterial strain *Sphingomonas* sp. 23L on fly ash. *Biol. Fret. Soil*, 45(4): 385-394.
- Izzo, A., M. Canright and T.D. Bruns, 2006. The effects of heat treatments on ectomycorrhizal resistant propagules and their ability to control bioassay seedlings. *Mycol. Res.*, 110(2): 196-202.
- Jonsson, L., A. Dahlberg, M.C. Nilsson, O. Zackrisson and O. Karen, 1999. Ectomycorrhizal fungal communities in late-successional Swedish boreal forests and their composition following wildfire. *Mol. Ecol.*, 8: 205-215.
- Kårén, O. and J.E. Nylund, 1997. Effect of ammonium sulphate on the community structure and biomass of ectomycorrhizal fungi in a Norway spruce stand in southwester Sweden. *Can. J. Bot.*, 75: 1628-1642.
- Kipfer, T., B. Moser, S. Egli, T. Wohlgemuth and J. Ghazoul, 2011. Ectomycorrhiza succession patterns in *Pinus sylvestris* forests after stand-replacing fire in the Central Alps. *Oecologia*, 167: 219-228.
- Klink, A., B. Letachowicz, J. Krawczyk and M. Wisłocka, 2006. The content of heavy metals in soil and silver birch leaves (*Betula pendula* Roth.) from Walbrzych and Gogol. *Polish J. Environ. Stud.*, 15: 347-350.

- Knoepp, J.D., L.F. DeBano and D.G. Neary, 2005. Soil Chemistry. In: Neary, D.G., K.C. Ryan and L.F. DeBano (Eds.), Wild Land Fire in Ecosystems: Effects of Fire on Soils and Water. General Technical Report. RMRS-GTR-42-vol. 4. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, pp: 63-71.
- Kopponen, P., M. Utriainen, K. Lukkari, S. Suntioinen, L. Karenlampi and S. Karenlampi, 2001. Clonal differences in copper and zinc tolerance of birch in metalsupplemented soils. Environ. Pollut., 112: 89-97.
- Koide, R.T., C. Fernandez and K. Petprakob, 2011. General principles in the community ecology of ectomycorrhizal fungi. Ann. For Sci., 68(1): 45-55.
- Kozdrój, J., Z. Piotrowska-Seget and P. Krupa, 2007. Mycorrhizal fungi and ectomycorrhiza associated bacteria isolated from an industrial desert soil protect pine seedlings against Cd (II) impact. Ecotoxicology, 16: 449-456.
- Krznaric, E., N. Verbruggen, J.H.L. Wevers, R. Carleer, J. Vangronsveld and J.V. Colpaert, 2009. Cd-tolerant *Suillus luteus*: A fungal insurance for pines exposed to Cd. Environ. Pol., 157:1581-1588.
- Leski, T., M. Rudawska and B. Kieliszewska-Ruziicka, 1995. Intraspecific aluminum response in *Suillus luteus* (L.) S.F. Gray., an ECM symbiotic of Scots Pine. Acta Societatis Botanic Rum Polonaise, 64: 97-105.
- Leyval, C., K. Turnau and K. Haselwandter, 1997. Effect of heavy metal pollution on mycorrhizal colonization and function: Physiological, ecological and applied aspects. Mycorrhiza, 7: 139-153.
- Lilleskov, E.A., T.J. Fahey and G.M. Lovett, 2001. Ectomycorrhizal fungal aboveground community change over an atmospheric nitrogen deposition gradient. Ecol. Appl., 11: 397-410.
- Lilleskov, E.A., T.J. Fahey, T.R. Horton and G.R. Lovett, 2002a. Belowground ectomycorrhizal fungal community change over a nitrogen deposition gradient in Alaska. Ecology, 83(1): 104-115.
- Lilleskov, E.A., E.A. Hobbie and T.J. Fahey, 2002b. Ectomycorrhizal fungal taxa differing in response to nitrogen deposition also differ in pure culture organic nitrogen use and natural abundance of nitrogen isotopes. New Phytol., 154: 219-231.
- Lingua, G., C. Franchin, V. Todeschini, S. Castiglione, S. Biondi, B. Burlando, V. Parravicini, P. Torrigiani and G. Berta, 2008. Arbuscular mycorrhizal fungi differentially affect the response to high zinc concentrations of two registered poplar clones. Environ. Pollut., 153:137-147.
- Ma, D.L., G. Yang and L. Mu, 2010. Morphological and molecular analyses of ectomycorrhizal diversity in *Pinus densiflora* seedlings. Symbiosis, 51: 233-238.
- Mah, K., L.E. Tackaberry, K.N. Egger and H.B. Massicotte, 2001. The impacts of broadcast burning after clear-cutting on the diversity of ectomycorrhizal fungi associated with hybrid spruce seedlings in central British Columbia. Can. J. Forest Res., 31: 224-235.
- Maia, L.C., A.M. Yano and J.W. Kimbrough, 1996. Species of ascomycota forming ectomycorrhizae. Mycotaxon, 57: 371-390.
- Markkola, A.M., J.U. Ahonen, M. Roitto, R. Strommer and M. Hyvarinen, 2002. Shift in ectomycorrhizal community composition in Scots pine (*Pinus sylvestris* L.) seedling roots as a response to nickel deposition and removal of lichen cover. Environ. Pollut., 120: 797-803.
- Marschner, H. and B. Dell, 1994. Nutrient uptake in mycorrhizal symbiosis. Plant Soil, 159: 89-102.
- Meharg, A.A. and J.W.G. Cairney, 2000. Ectomycorrhizas-extending the capabilities of rhizosphere remediation? Soil Bio. Biochem., 32: 1475-1484.
- Mogge, B., C. Loferer, R. Agerer, P. Huzler and A. Hartman, 2000. Bacterial community structure and colonisation patterns of *Fagus sylvatica* L. ectomycorrhizospheres as determinate by fluorescence in situ hybridization and co focal laser scanning microscopy. Mycorrhiza, 9: 271-278.
- Nilsson, L.O., 2004. External mycelia of mycorrhizal fungi. Ph.D. Thesis, Department of Ecology, Lund University, Sweden.
- Nurmiaho-Lassila, E.L., S. Timonen, K. Haahtela and R. Sen, 1997. Bacterial colonization patterns of intact *Pinus Sylvester*'s mycorrhizospheres in dry pine forest soil: An electron microscopy study. Can. J. Microbial., 43:1017-1035.
- Palfner, G., 1997. *Descolea anarctica* Singer + *Nothofagus alpina* (Poepp. and Endl.), Oerst. Descry Entomic, 2: 7-12.
- Palfner, G., 2001. Taxonomist studied an ektomykorrhizen aus den nothofagus-waldern mittelsüdchiles: Bibliotheca mycological. J. Cramer. Berlin., 190: 243.
- Palfner, G., 2002. *Cortinarius Magellanicus*. In: Agerer R (Ed) Color Atlas of Ectomycorrhizae (1987–2006). Einhorn-Verlag Schwäbisch Gmünd, Germany, pp: 153.
- Palmer, J.M., D.L. Lindner and T.J. Volk, 2008. Ectomycorrhizal characterization of an American chestnut (*Castaneadentata*)-dominated community in western Wisconsin. Mycorrhiza, 19:27-36.
- Palfner, G., M.A. Casanova-Katny and D.J. Read, 2005. The mycorrhizal community in a forest chronosequence of Sitka spruce [*Picea sitchensis* (Bong.) Carr Northern England. Mycorrhiza, 15: 571-579.
- Palfner, G., M.I. Canseco and A. Casanova-Katny, 2008. Post-fire seedlings of *Nothofagus alpina* in Southern Chile show strong dominance of a single ectomycorrhizal fungus and a vertical shift in root architecture. Plant Soil, 313: 237-250.

- Parrent, J.L., W.F. Morris and R. Vilgalys, 2006. CO<sub>2</sub>-enrichment and nutrient availability alter ectomycorrhizal fungal communities. *Ecology*, 87: 2278-2287.
- Parrent, J.L. and R. Vilgalys, 2007. Biomass and compositional responses of ectomycorrhizal fungal hyphae to elevated CO<sub>2</sub> and nitrogen fertilization. *New Phytol.*, 176:164-174.
- Peay, K.G., T.D. Bruns, P.G. Kennedy, S.E. Bergemann and M. Garbelotto, 2007. A strong species-area relationship for eukaryotic soil microbes: Island size matters for ectomycorrhizal fungi. *Ecol. Lett.*, 10: 470-480.
- Peter, M., F. Ayer and S. Egli, 2001a. Nitrogen addition in a Norway spruce stand altered macromycetes sporocarp production and belowground ectomycorrhizal species composition. *New Phytol.*, 149: 311-325.
- Peter, M., F. Ayer, S. Egli and R. Honegger, 2001b. Above and below-ground community of ectomycorrhizal fungi in three Norway spruce (*Picea abies*) stands in Switzerland. *Can. J. Bot.*, 79: 1134-1151.
- Petersen, P.M., 1970. Danish fireplace fungi: An ecological investigation of fungi on burns. *Dansk Bot. Arkiv.*, 27: 1-97.
- Poole, E.J., G.D. Bending, J.M. Whipps and D.J. Read, 2001. Bacteria associated with *Pinus sylvestris-Lactarius rufus* ectomycorrhizas and their effects on mycorrhiza formation in vitro. *New Phytol.*, 151: 743-751.
- Wang, Q., X.H. He and L.D. Guo, 2012. Ectomycorrhizal fungus communities of *Quercus liaotungensis* Koidz of different ages in a northern China temperate forest. *Mycorrhiza*, 22: 461-470.
- Regvar, M., K. Vogel-Mikuš, N. Kugonič, B. Turk and F. Baltič, 2006. Vegetation and mycorrhizal successions at a metal polluted site: Indications for the direction of phytostabilisation? *Environ. Pollut.*, 144: 976-984.
- Richard, F., P.A. Moreau, M.A. Selosse and M. Gardes, 2004. Diversity and fruiting patterns of ectomycorrhizal and saprobe fungi in an old growth Mediterranean forest dominated by *Quercus ilex* L. *Can. J. Bot.*, 82: 1711-1729.
- Richard, F., S. Millot, M. Gardes and M.A. Selosse, 2005. Diversity and specificity of ectomycorrhizal fungi retrieved from an old-growth Mediterranean forest dominated by Quercus ilex. *New Phytol.*, 166: 1011-1023.
- Román, M. and A.M. Miguel, 2005. Post-fire, seasonal and annual dynamics of the ectomycorrhizal community in a Quercus ilex L. forest over a 3-year period. *Mycorrhiza*, 15: 471-482.
- Rygiewicz, P.T., M.G. Johnson, L.M. Ganio, D.T. Tingey and M.J. Storm, 1997. Lifetime and temporal occurrence of ectomycorrhizae on ponderosa pine (*Pinus ponderosa* Laws.) seedlings grown under varied atmospheric CO<sub>2</sub> and nitrogen levels. *Plant Soil*, 189: 275-287.
- Sakakibara, S.M., M.D. Jones, M. Gillespie, S.M. Hagerman, M.E. Forrest, S.W. Simard and D.M. Durall, 2002. A comparison of ectomycorrhiza identification based on morphotyping and PCR-RFLP analysis. *Mycol. Res.*, 106: 868-878.
- Sell, J., A. Kayser, R. Schulin and I. Brunner, 2005. Contribution of ectomycorrhizal fungi to cadmium uptake of poplars and willows from a heavily polluted soil. *Plant Soil*, 277: 245-253.
- Shishido, M., H.B. Massicotte and C.P. Chanway, 1996. Effect of plant growth promoting Bacillus strains on pine and spruce seedling growth and mycorrhizal infection. *Ann. Bot.*, 77: 433-441.
- Smith, S.E. and D.J. Read, 1997. *Mycorrhizal Symbiosis*. Academic, San Diego.
- Stendell, E.R., T.R. Horton and T.D. Bruns, 1999. Early effects of prescribed fire on the structure of the ectomycorrhizal fungus community in a Sierra Nevada ponderosa pine forest. *Mycol. Res.*, 103(10): 1353-1359.
- States, J.S. and W.S. Gaud, 1997. Ecology of hypogeous fungi associated with ponderosa pine. I. patterns of distribution and sporocarp production in some Arizona forests. *Mycological*, 89: 712-726.
- Taylor, A.F.S., 2002. Fungal diversity in ectomycorrhizal communities: Sampling effort and species detection. *Plant Soil*, 244(1): 19-28.
- Tedersoo, L., U. Koljalg, N. Hallenberg and K.H. Larsson, 2003. Fine scale distribution of ectomycorrhizal fungi and roots across substrate layers including coarse woody debris in a mixed forest. *New Phytol.*, 159:153-165.
- Torres, P. and M. Honrubia, 1997. Changes and effects of a natural fire on ectomycorrhizal inoculum potential of soil in a Pinus halepensis. *Forest Ecol. Manage.*, 96:189-196.
- Treseder, K.K., 2004. A meta-analysis of mycorrhizal responses to nitrogen, phosphorus and atmospheric CO<sub>2</sub> in field studies. *New Phytol.*, 164: 347-355.
- Trowbridge, J. and A. Jumpponen, 2004. Fungal colonization of shrub willow roots at the forefront of a receding glacier. *Mycorrhiza*, 14: 283-293
- Tuininga, A.R. and J. Dighton, 2004. Changes in ectomycorrhizal communities and nutrient availability following prescribed burns in two upland pine-oak forests in the New Jersey Pine Barrens. *Can. J. Res.*, 34(8):1755-1765.
- Twieg, B.D., D.M. Durall and S.W. Simard, 2007. Ectomycorrhizal fungal succession in mixed temperate forests. *New Phytol.*, 176: 437-447.

- Twieg, B.D., D.M. Durall, S.W. Simard and M.D. Jones, 2009. Influence of soil nutrients on ectomycorrhizal communities in a chronosequence of mixed temperate forests. *Mycorrhiza*, 19: 305-316.
- Vallino, M., N. Massa, E. Lumini, V. Bianciotto, G. Berta and P. Bonfante, 2006. Assessment of carbuncular mycorrhizal fungal diversity in roots of *Solidago gigantea* growing in a polluted soil in Northern Italy. *Environ. Microbiol.*, 8: 971-983.
- Visser, S., 1995. Ectomycorrhizal fungal succession in jack pine stands following wildfire. *New Phytol.*, 129: 389-401.
- Vrålstad, T., A. Holst-Jensen and T. Schumacher, 1998. The postfire discomycete *Geopyxis carbonaria* (Ascomycota) is a biotrophic root associate with Norway spruce (*Picea abies*) in nature. *Mol. Ecol.*, 7: 609-616.
- Walker, R.F., D.R. Geisinger, D.W. Johnson and J.T. Ball, 1995. Interactive effects of atmospheric CO<sub>2</sub> enrichment and soil N on growth and ectomycorrhizal colonization of *ponderosa pine* seedlings. *Forest Sci.*, 41: 491-500.
- Walker, J.F., O.K. Miller and J.L. Horton, 2008. Seasonal dynamics of ectomycorrhizal fungus assemblages on oak seedlings in the southeastern appalachian mountains. *Mycorrhiza*, 18: 123-132.
- Warcup, J.H., 1990. Occurrence of ectomycorrhizal and saprophytic discomycetes after a wild fire in a eucalypt forest. *Mycol. Res.*, 94:1065-1069.
- Wiklund, K., L.O. Nilsson and S. Jacobson, 1995. Effect of irrigation, fertilization and artificial drought on basidioma production in a Norway spruce stand. *Can. J. Bot.*, 73: 200-208.
- Wurzburger, N, M.I. Bidartondo and C.S. Bledsoe, 2001. Characterization of Pinus ectomycorrhizas from conifer and pygmy forest using morph typing and molecular methods. *Can. J. Bot.*, 79:1211-1216.
- Yamada, A. and K. Katsuya, 1995. Mycorrhizal association of isolates with *Pinus densiflora* seedlings. *Mycoscience*, 36: 315-323.