

Research Article

Microbial Involvement in Growth of Paddy

¹Febri Doni, ¹Najeeb Kaid Nasser Al-Shorgani, ¹El Mubarak Musa Tibin, ²Nawal Noureldaim Abuelhassan, ³Anizan Isahak, ¹Che Radziah Che Mohd. Zain and ¹Wan Mohtar Wan Yusoff

¹Faculty Science and Technology, School of Biosciences and Biotechnology,

²Faculty Sciences and Technology, School of Chemical Sciences and Food Technology,

³Faculty Sciences and Technology, School of Environmental and Natural Resources Sciences, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

Abstract: This study reviews the role of Plant Growth Promoting Rhizobacteria (PGPR), Plant Growth Promoting Fungi (PGPF) and Mycorrhizal Fungi in the enhancement of rice growth and yield. A total 70 papers spanning from 1970 to 2013 was screened and noted its conclusion to enhance rice growth and yield. The mechanisms employed by microbes in enhancing rice growth and yield includes; the production of growth regulating substances, phosphate solubilization, N-fixation, cellulose degradation and siderophore production. Some microbes are also involved in cell regulation and signaling in rice plants. We can conclude that there are significant influences on the growth and yield of rice plant by microbes.

Keywords: Growth PGPR, mycorrhizal fungi, PGPF, rice

INTRODUCTION

Currently growth of paddy uses a lot of chemical fertilizer and chemical pesticide. Over the years this has led to serious environmental problems such as depletion of soil quality and health, emergence of resistant pathogens and elimination of soil microbes involvement in growth of paddy. The increase of production of paddy must be achieved through improvement in agricultural productivity. Microbes considered as beneficial are key factor in maintaining soil quality and paddy production. As such decades, Interest in beneficial microorganism in rice has increased due to their potential use as plant growth regulator (Diem *et al.*, 1978; Barraquio *et al.*, 1986; Mehnaz *et al.*, 2000; Jha and Subramanian, 2012).

Previous studies have reviewed the microbial producers of plant growth stimulators and their practical use (Tsavkelova *et al.*, 2005), use of plant growth-promoting bacteria for biocontrol of plant diseases (Compant *et al.*, 2005), plant-microbes interactions in enhanced fertilizer-use efficiency (Adesemoye and Kloepper, 2009), bacterial and fungal contributions to carbon sequestration in agroecosystems (Six *et al.*, 2006), biological nitrogen fixation (Kumar and Rao, 2012) and impact of root exudates and plant defense signaling on bacterial communities in the rhizosphere (Doornbos *et al.*, 2012).

This review discusses the ability of Plant Growth Promoting Rhizobacteria (PGPR) to enhance rice

growth and yield, the ability of Plant Growth Promoting Fungi (PGPF) to enhance rice growth and yield, the ability of mycorrhizal fungi to enhance rice growth and yield and rice plant and microorganism molecular interaction.

RESULTS AND DISCUSSION

Plant growth promoting rhizobacteria: PGPR (Plant Growth Promoting Rhizobacteria) are the rhizosphere bacteria which may grow in, on or around plant tissues and have ability to improve plant growth by several mechanisms namely Phosphate-solubilizing, siderophore-producing, nitrogen-fixing, phytohormone-producing and exhibiting antifungal activity. (Bhattacharyya and Jha, 2012; Vessey, 2003). The mechanism of PGPR to improve plant growth is highlighted on Table 1.

Reports on the application of PGPR to improve rice growth and yield have been published by many researchers. Carreres *et al.* (1996) stated that application of cyanobacteria to the rice plant significantly increased the uptake of N by the rice plant. Further, Malik *et al.* (1997) stated that five strains of bacteria, namely, *Azospirillum lipoferum* N-4, *A. brasilense* Wb-3, *Pseudomonas* 96-51, *Zoogloea* Ky-1 and *Azoarcus* K-1 were able to increase biomass and nitrogen content of the rice tissue. Ashrafuzzaman *et al.* (2009) isolated 10 PGPR strains from Mymensingh soil to improve rice growth. Most of isolates resulted in a

Corresponding Author: Febri Doni, Faculty Science and Technology, School of Biosciences and Biotechnology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

This work is licensed under a Creative Commons Attribution 4.0 International License (URL: <http://creativecommons.org/licenses/by/4.0/>).

Table 1: Growth factor produced by PGPR

PGPR	Factor produced	References
<i>Azospirillum lipoferum</i>	Gibberellins N-Fixation	Cassan <i>et al.</i> (2001) and Nayak <i>et al.</i> (1986)
<i>Azospirillum brasilense</i>	Gibberellins IAA-N-Fixation	Cassan <i>et al.</i> (2001), Mehnaz and Lazarovits (2006) and Tien <i>et al.</i> (1976)
Zoogloea	N-Fixation	Xie and Yokota (2006)
Azoarcus	N-Fixation	Hurek <i>et al.</i> (2012)
Bacillus	Phosphate-solubilizing	Rodríguez and Fraga (1999)
Rhizobium	N-Fixation	Yani <i>et al.</i> (2000)
<i>Pseudomonas fluorescens</i>	N-Fixation Phosphate-solubilizing Siderophore producing	Park <i>et al.</i> (2004) and Vyas and Gulati (2009)
<i>Pseudomonas putida</i>	Siderophore producing Phosphate-solubilizing	Kloepper <i>et al.</i> (1980)
Azetobacter	N-Fixation	Kloepper <i>et al.</i> (1980) and Wahyudi <i>et al.</i> (2011)
Azorhizobium	N-Fixation	Park <i>et al.</i> (2004)
Azospirillum	N-Fixation	Anyia <i>et al.</i> (2004)
		Park <i>et al.</i> (2004)

Table 2: Growth factor produced by PGPF

PGPF	Factor produced	References
<i>Gliocladium virens</i>	Lytic enzymes	Sreenivasaprasad and Manibhushanrao (1990)
<i>Trichoderma virens</i>	Auxin Jalicylic acid Jasmonic acid	Contreras-Cornejo <i>et al.</i> (2009, 2011)
<i>Trichoderma harzianum</i>	Phosphate-solubilizing Siderophore producing	Saravanakumar <i>et al.</i> (2013) and Rawat and Tewari (2011)
<i>Trichoderma viride</i>	Phosphate-solubilizing Siderophore producing cellulose degrading	Saravanakumar <i>et al.</i> (2013), Rawat and Tewari (2011) and Jiang <i>et al.</i> (2011)
<i>Aspergillus niger</i>	IAA Gibberellins Phytase producing	Bilkay <i>et al.</i> (2010) and Gujar <i>et al.</i> (2013)

significant increase in plant height, root length and dry matter production of shoot and root of rice seedlings. Mia *et al.* (2012) has inoculated 4 PGPR strains to rice seedlings growth. The results stated that inoculation those PGPR strains have been increasing the seedling emergence, seedling vigor and root growth.

The ability of PGPR to enhance rice yield has been reported in several studies. Sakhivel and Gnanamanickam (1987) reported that application of *Pseudomonas fluorescens* to rice plant significantly improve plant height, number of tillers and grain yields. Mohammadinejad-Babandeh *et al.* (2012) demonstrated that Azotobacter, Azorhizobium and Azospirillum were able to enhance rice yield components. Vahed *et al.* (2012) found that Phosphate Solubilizing Bacteria (PSB) had a significant influence on grain yield and biological yield of rice. Gopalakrishnan *et al.* (2012) reported that four PGPR isolates from rhizosphere of a SRI field significantly enhanced rice tiller numbers, stover and grain yields, total dry matter, root length, volume and dry weight. Ruíz-Sánchez *et al.* (2011) examined the response of rice plants to inoculation with *A. brasilense*. Result showed that *A. brasilense* is able to enhance ascorbate content on rice plants. Pedraza *et al.* (2009) assessed the Azospirillum inoculation on effect on grain yield in a rice rainfed crop. The results suggested that Azospirillum significantly increased rice yield. Further, Banayo *et al.* (2012) evaluated three different biofertilizers. The result showed significant yield increases for all biofertilizers in some seasons but the most consistent results were achieved by the *Azospirillum*-based biofertilizer.

Plant growth promoting fungi: Plant Growth Promoting Fungi (PGPF) is a class of beneficial fungi that have ability to enhance plant growth. The PGPF are including; hormone production, substrate degradation

(mineralization) and suppression of deleterious microorganism (Hyakumachi, 1994; Newwani *et al.*, 2010). The mechanism of plant growth promotion by PGPF is listed in Table 2.

The ability of PGPF to enhance growth and yield of rice has been reported by many researchers. Four fungal (*Gliocladium virens*, *Trichoderma virens*, *T. harzianum* and *Aspergillus niger*) were examined for their effect on germination, root and shoot length and seedling weight of rice. The results suggested that seed root length, shoot height and fresh weight of rice seedlings were significant increased (Mishra and Sinha, 2000). Further, Al-Taweil *et al.* (2009) has applied *T. viride* to rice seedling and the results reported that *T. viride* is effectively increasing rice seedlings growth. Khan *et al.* (2005) examined activity of *T. harzianum* on rice seed germination and seedling vigor. The result showed that *T. harzianum* has ability to increase seedling emergence, root and shoot length, fresh and dry weight of root of rice seedlings. The same result of the ability of *T. harzianum* on improving rice seedling growth was also reported by Shukla *et al.* (2012). Choi *et al.* (2010) reported that imidazole-4-carboxamide (ICA), a plant-growth regulating compound that was isolated from a fairy ring forming fungus, *Lepista sordida*, in a greenhouse experiment, this compound increased rice grain yield by 26% compared with control. Amprayn *et al.* (2012) investigated the ability of *Candidatropicalis* HY (CtHY) to stimulate rice seedlings growth. The application of CtHY on germinated seedlings resulted in better rice plant root growth and the colonization of CtHY was confirmed to persist on plant roots at least for 3 weeks. Furthermore, Banaay *et al.* (2012) reported that *T. ghanense* has ability to promote seedlings growth of aerobic rice variety.

Mycorrhizal fungi: Arbuscular mycorrhizal (AM) fungi are group of beneficial fungi that symbioses with

the roots of around 80% of vascular plants and often increasing Phosphate (P) uptake and growth (Smith *et al.*, 2003). Hajiboland *et al.* (2009) stated that Mycorrhizal colonization on rice significantly contributes to the uptake of P and K in rice plants. The ability of AM to enhance rice growth and yield has been reported by Secilia and Bagyaraj (1992), Secilia and Bagyaraj (1994), Solaiman and Hirata (1997), Li *et al.* (2011) and Zhang *et al.* (2012). Yeasmin *et al.* (2007) stated that Mycorrhizal fungi significantly improved rice plants growth by increasing soil nutrients such as nitrogen and phosphorus. Isahak *et al.* (2012) examined the influence of AM to influence rice seedlings growth, as the result showed that AM is significantly enhancing plant height. Zhang *et al.* (2005) investigated the ability of AM to increase upland rice growth under combined soil contamination. The results showed that Mycorrhizal colonization on upland rice had a large influence on rice growth by increasing the shoot and root biomass. The inoculation of AM also gave the protective effects on upland rice under the combined soil contamination Further, Xu *et al.* (2013) stated that AM is able on enhancing rice production when growing in As-contaminated soils.

Paddy and microorganism molecular interaction:

Plant-microbe interactions may occur at several core spaces, specific spaces and sub-spaces within the specific space of choice and inter-spaces. These may include amongst others; Phyllosphere, Endosphere and Rhizosphere as core spaces. Phyllosphere is related with the aerial domains of the plants and endosphere being related with the internal cellular systems and its attendant transport systems. Rhizosphere can be defined as any volume of soil specially and specifically influenced by the plant roots or in association with the roots and plant-produced material. Rhizosphere, the term, can be defined as any volume of soil specially influenced by the plant roots or in association with the roots and plant-produced material. Bhattacharyya and Jha (2012) also mentioned in relation to the existence of the secorespaces. Plantexudes some phenolic compounds into the rhizosphere as signal to attract microorganism (Peters and Verma, 1990; Bais *et al.*, 2004). Bacilio-Jiminez *et al.* (2003) identified rice root exudates which fall into two separate groups, amino acids (histidine, proline, valine, alanine, glycine, aspartic, arginine, tyrosine and methionine) and carbohydrate (mannose, galatose, glucose and glucuronic acid).

The abundance of rice root exudates might attract microorganism to colonize rice root that penetrate to root tissue. Reinhold-Hurek and Hurek (1997) examined the ability of *Azoarcus* to colonize rice roots endophytically. The result showed that bacteria invade the roots in the zone of elongation and differentiation, colonize the cortex intra- and inter-cellularly and

penetrate deeply into the vascular system, entering xylem vessels, allowing systemic spreading into the rice shoot. Blilou *et al.* (2000) stated that the expression of a Lipid Transfer Protein (LTP) gene is regulated in *Oryza sativa* roots in response to colonization by the mycorrhizal fungus *Glomus mosseae*. And then, Transcript levels increased when the fungus forms appressoria and penetrate the root epidermis and decreased at the onset of the intercellular colonization of the root cortex. Further, Rediers *et al.* (2003) examined *Pseudomonas stutzeri* A15 genes that are switched on during rice root colonization and are switched off during free-living growth on synthetic medium. This strain is able to promote rice growth. Further, Guimil *et al.* (2005) stated 12 genes has transcribed only when the mycorrhizal fungus *Glomus intraradices* colonized rice root and those 12 genes were not detected in the absence of symbiosis.

CONCLUSION

This review has shown that there are significant influences of microbes to the growth and yield of rice plant. Many studies investigate the role of microbes to enhance rice growth and yield including producing of growth regulating substances, phosphate-solubilizing, N-fixation, cellulose degrading and siderophore producing. Some microbes are also involving in cell regulation and signaling in rice plants.

ACKNOWLEDGMENT

This study was funded by Universiti Kebangsaan Malaysia under grant Komuniti-2012-001.

REFERENCES

- Adesemoye, A.O. and J.W. Kloepper, 2009. Plant-microbes interactions in enhanced fertilizer-use efficiency. *Appl. Microbiol. Biot.*, 85: 1-12.
- Al-Taweil, H.I.B., M. Osman, A.A. Hamid and W.M. Wan Yusoff, 2009. Development of microbial inoculants and the impact of soil application on rice seedlings growth. *Am. J. Agr. Biol. Sci.*, 4: 79-82.
- Amprayn, K.O., T.R. Michael, M. Kecskés, L. Pereg, H.T. Nguyen and I.R. Kennedy, 2012. Plant growth promoting characteristics of soil yeast (*Candida tropicalis* HY) and its effectiveness for promoting rice growth. *Appl. Soil Ecol.*, 61: 295-299.
- Anyia, A.O., D.J. Archambault and J.J. Slaski, 2004. Growth promoting effects of the diazotroph azorhizobium caulnodans on canadian wheat cultivars. *Proceedings of the 4th International Crop Science Congress. Brisbane, Australia, 26 September-1 October.*

- Ashrafuzzaman, M., F.A. Hossen, M.R. Ismail, M.A. Hoque, M.Z. Islam, S.M. Shahidullah and S. Meon, 2009. Efficiency of Plant Growth-Promoting Rhizobacteria (PGPR) for the enhancement of rice growth. *Afr. J. Biotechnol.*, 8: 1247-1252.
- Bacilio-Jimenez, S., S. Aguilar-Flores, E. Ventura-Zapata, E. Perez-Campos, S. Bouquelet and E. Zenteno. 2003. Chemical characterization of root exudates from rice (*Oryza Sativa* L.) and their effects on chemotactic response of endophytic bacteria. *Plant Soil*, 249: 271-277.
- Bais, H.P., S.W. Park, T.L. Weir, R.M. Callaway and J.M. Vivanco, 2004. How plants communicate using the underground information superhighway. *Trends Plant Sci.*, 9: 26-32.
- Banaay, C.G.B., V.C. Cuevas and C.M. Vera Cruz, 2012. *Trichoderma ghanense* promotes plant growth and controls disease caused by *pythium arrhenomanes* in seedlings of aerobic rice variety apo. College Agriculture, University of Philippines, Los Banos, Laguna 4031, Philippines, ISSN: 0031-7454.
- Banayo, N.P.M., P.C.S. Cruz, E.A. Aguilar, R.B. Badayos and S.M. Haefele, 2012. Evaluation of bio-fertilizers in irrigated rice: Effects on grain yield at different fertilizer rates. *Agriculture*, 2: 73-86.
- Barraquio, W.L., J.K. Ladha, H.Q. Yao and I. Watanabe, 1986. Antigenic relationship of nitrogen-fixing pseudomonas strain H8 to various known cultures and rice rhizosphere isolates studied by indirect enzyme-linked immunosorbent assay (ELISA). *Can. J. Microbiol.*, 32: 402-408.
- Bhattacharyya, P.N. and D.K. Jha, 2012. Plant Growth-Promoting Rhizobacteria (PGPR): Emergence in agriculture. *World J. Microbiol. Biotechnol.*, 28: 1327-1350.
- Bilkay, I.S., Ş. Karakoc and N. Aksoz, 2010. Indole-3-acetic acid and gibberellic acid production in *Aspergillus niger*. *Turk. J. Biol.*, 34: 313-318.
- Blilou, I., J.A. Ocampo and J.M. García-Garrido, 2000. Induction of *Ltp* (*Lipid Transfer Protein*) and *Pal* (*Phenylalanine ammonia-lyase*) gene expression in rice roots colonized by the arbuscular mycorrhizal fungus *Glomus mosseae*. *J. Exp. Bot.*, 51: 1969-1977.
- Carreres, R. González Tomé, J. Sendra, R. Ballesteros, E. Fernández Valiente, A. Quesada, M. Nieva and F. Leganés, 1996. Effect of nitrogen rates on rice growth and biological nitrogen fixation. *J. Agric. Sci.*, 127: 295-302.
- Cassan, F., R. Bottini, G. Schneider and P. Piccoli, 2001. *Azospirillum brasilense* and *Azospirillum lipoferum* hydrolyze conjugates of GA20 and metabolize the resultant aglycones to GA1 in seedlings of rice dwarf mutants. *Plant Physiol.*, 125: 2053-2058.
- Choi, J.H., N. Abe, H. Tanaka, K. Fushimi, Y. Nishina, A. Morita, Y. Kirriwa, R. Motohashi, D. Hashizume, H. Koshino and H. Kawagishi, 2010. Plant-growth regulator, imidazole-4-carboxamide, produced by the fairy ring forming fungus *Lepista sordida*. *J. Agr. Food Chem.*, 58: 9956-9959.
- Compant, S., B. Duffy, J. Nowak, C. Clement and E.A. Barka, 2005. Use of plant growth-promoting bacteria for biocontrol of plant diseases: Principles, mechanisms of action and future prospects. *Appl. Environ. Microbiol.*, 71: 4951-4959.
- Contreras-Cornejo, H.A., L. Macías-Rodríguez, C. Cortés-Penagos and J. López-Bucio, 2009. *Trichoderma virens*: A plant beneficial fungus enhances biomass production and promotes lateral root growth through an auxin-dependent mechanism in arabidopsis. *Plant Physiol.*, 149: 1579-1592.
- Contreras-Cornejo, H.A., L. Macías-Rodríguez, E. Beltrán-Peña, A. Herrera-Estrella and J. López-Bucio, 2011. *Trichoderma*-induced plant immunity likely involves both hormonal- and camalexin-dependent mechanisms in *Arabidopsis thaliana* and confers resistance against necrotrophic fungus *Botrytis cinerea*. *Plant Signal Behav.*, 6: 1554-1563.
- Diem, G., M. Rougier, I. Hamad-Fares, J.P. Balandreau and Y.R. Dommergues, 1978. Colonization of rice roots by diazotroph bacteria. Environmental role of nitrogen-fixing blue-green algae and symbiotic bacteria. In: Granhall, U. (Ed.), *Ecol. Bull.* (Stockholm), 26: 305-311.
- Doornbos, R.F., L.C. van Loon and P.A.H.M. Bakker, 2012. Impact of root exudates and plant defense signaling on bacterial communities in the rhizosphere. A review. *Agron. Sust. Dev.*, 32: 227-243.
- Gopalakrishnan, S., H.D. Upadhyaya, S. Vadlamudi, P. Humayun, M.S. Vidya, G. Alekhya, A. Singh, R. Vijayabharathi, R.K. Bhimineni, M. Seema, A. Rathore and O. Rupela, 2012. Plant growth-promoting traits of biocontrol potential bacteria isolated from rice rhizosphere. *Springer Plus*, 1: 71, DOI: 10.1186/2193-1801-1-71.
- Guimil, S., H.S. Chang, T. Zhu, A. Sesma, A. Osbourn, C. Roux, V. Ioannidis, E.J. Oakeley, M. Docquier, P. Descombes, S.P. Briggs and U. Paszkowski, 2005. Comparative transcriptomics of rice reveals an ancient pattern of response to microbial colonization. *Proc. Natl. Acad. Sci. USA.*, 102: 8066-8070.
- Gujar, P.D., K.P. Bhavsar and J.M. Khire, 2013. Effect of phytase from *Aspergillus niger* on plant growth and mineral assimilation in wheat (*Triticum aestivum* Linn.) and its potential for use as a soil amendment. *J. Sci. Food Agric.*, 93(9): 2242-7.

- Hajiboland, R., N. Aliasghar zad and R. Barzeghar, 2009. Phosphorus mobilization and uptake in mycorrhizal rice (*Oryza sativa* L.) plants under flooded and non-flooded conditions. *Acta Agric. Slov.*, 93: 153-161.
- Hurek, T., L.L. Handley, B. Reinhold-Hurek and Y. Piché, 2012. Azoarcus grass endophytes contribute fixed nitrogen to the plant in an unculturable state. *Mol. Plant Microb. Interact.*, 15: 233-242.
- Hyakumachi, M., 1994. Plant growth fungi from turf grass rhizosphere with potential for disease suppression. *Soil Microorgan.*, 44: 53-68.
- Isahak, A., A. Ahmad, A.B. Rosenani and H. Jamil, 2012. SRI rice crop establishment. *Trans. Malaysian Soc. Plant Physiol.*, 20: 20-20.
- Jha, Y. and R.B. Subramanian, 2012. Isolation of root associated bacteria from the local variety of rice GJ-. *World Res. J. Geoinform.*, 1: 21-26.
- Jiang, X., A. Geng, N. He and Q. Li, 2011. New isolate of *Trichoderma viride* strain for enhanced cellulolytic enzyme complex production. *J. Biosci. Bioeng.*, 111: 121-127.
- Khan, A.A., A.P. Sinha and Y.P.S. Rathi, 2005. Plant growth promoting activity of *Trichoderma herzianum* on rice seed germination. *Indian J. Agric. Res.*, 39: 256-262.
- Klopper, J.W., J. Leong, M. Teintze and M.N. Schroth, 1980. Enhanced plant growth by siderophores produced by plant growth-promoting rhizobacteria. *Nature*, 286: 885-886.
- Kumar, S.R.S. and K.V.B. Rao, 2012. Biological nitrogen fixation: A review. *Int. J. Adv. Life Sci.*, 1: 1-6.
- Li, W., M. Fang, Z. Shujuan and Z. Xue, 2011. Effect of glomus mosseae inoculation on growth and reproduction of rice. *Proceeding of the International Conference on Information Technology and Agricultural Engineering (ICITAE 2011)*. Sanya, China.
- Malik, K.A., B. Rakhshanda, M. Samina, G. Rasul, M.S. Mirza and S. Ali, 1997. Association of nitrogen-fixing, plant-growth-promoting rhizobacteria (PGPR) with kallar grass and rice. *Plant Soil*, 194: 37-44.
- Mehnaz, S. and G. Lazarovits, 2006. Inoculation effects of *Pseudomonas putida*, *Gluconacetobacter azotocaptans* and *Azospirillum lipoferum* on corn plant growth under greenhouse conditions. *Microb. Ecol.*, 51: 326-35.
- Mehnaz, S., M. Sajjad Mirza, H. Jacqueline, B. Rene, N. Philippe, B. Asghari and A.M. Kauser, 2000. Isolation and 16S rRNA sequence analysis of the beneficial bacteria from the rhizosphere of rice. *Can. J. Microbiol.*, 4: 110-117.
- Mia, M.A.B., Z.H. Shamsuddin and M. Mahmood, 2012. Effects of rhizobia and plant growth promoting bacteria inoculation on germination and seedling vigor of lowland rice. *Afr. J. Biotechnol.*, 11: 3758-3765.
- Mishra, D.S. and A.P. Sinha, 2000. Plant growth-promoting activity of some fungal and bacterial agents on rice seed germination and seedling growth. *Trop. Agric.*, 77: 188-191.
- Mohammadinejad-Babandeh, S.N., H.R. Doroodian and H. Besharati, 2012. Effect of bio-bacterial (*Azotobacter*, *Azorhizobium*, *Azospirillum*) on yield and yield components of rice in Bandar-Anzali, North of Iran. *Res. J. Biol. Sci.*, 7: 244-249.
- Nayak, D.N., J.K. Ladha and I. Watanabe, 1986. The fate of marker *Azospirillum lipoferum* inoculated into rice and its effect on growth, yield and N₂ fixation of plants studied by acetylene reduction, ¹⁵N₂ feeding and ¹⁵N dilution techniques. *Biol. Fertil. Soils*, 1: 7-14.
- Neuwani, V., P. Doshi, T. Saha and S. Rajkumar, 2010. Isolation and characterization of a fungal isolate for phosphate solubilization and plant growth promoting activity. *J. Yeast Fungal Res.*, 1: 009-014.
- Park, M., C. Kim, J. Yang, H. Lee, W. Shin, S. Kim and T. Sa, 2004. Isolation and characterization of diazotrophic growth promoting bacteria from rhizosphere of agricultural crops of Korea. *J. Microsc.*, 160: 127-133.
- Pedraza, R.O., C.H. Bellone, S.C. de Bellone, P.M.F.B. Sorte and K.R. dos Santos Teixeira, 2009. *Azospirillum* inoculation and nitrogen fertilization effect on grain yield and on the diversity of endophytic bacteria in the phyllosphere of rice rainfed crop. *Eur. J. Soil Biol.*, 45: 36-43.
- Peters, N.K. and D.P.S. Verma, 1990. Phenolic compounds as regulators of gene expression in plant-microbe interactions. *Mol. Plant-Microb. Interact.*, 3: 84-88.
- Rawat, R. and L. Tewari, 2011. Effect of abiotic stress on phosphate solubilization by biocontrol fungus *Trichoderma sp.* *Curr. Microbiol.*, 62: 1521-1526.
- Rediers, H., V. Bonnacerrere, P.B. Rainey, K. Hamonts, J. Vanderleyden and R. De Mo, 2003. Development and application of a DapB-based in vivo expression technology system to study colonization of rice by the endophytic nitrogen-fixing bacterium *pseudomonas stutzeri* A15. *Appl. Environ. Microbiol.*, 69: 6864-6874.
- Reinhold-Hurek, B. and T. Hurek, 1997. *Azoarcus* spp. and their interactions with grass roots. *Plant Soil*, 194: 57-64.
- Rodríguez, H. and R. Fraga, 1999. Phosphate solubilizing bacteria and their role in plant growth promotion. *Biotechnol. Adv.*, 17: 319-339.
- Ruíz-Sánchez, M., E. Armada, Y. Munoz, I.E.G. de Salamone, R. Aroca, J.M. Ruíz-Lozano and R. Azcón, 2011. *Azospirillum* and arbuscular mycorrhizal colonization enhance rice growth and physiological traits under well-watered and drought conditions. *J. Plant Physiol.*, 168: 1031-1037.

- Sakthivel, N. and S.S. Gnanamanickam, 1987. Evaluation of *Pseudomonas fluorescens* for suppression of sheath rot disease and for enhancement of grain yields in rice (*Oryza sativa* L.). *Appl. Environ. Microbiol.*, 53: 2056-2059.
- Saravanakumar, K., V.S. Arasu and K. Kathiresan, 2013. Effect of trichoderma on soil phosphate solubilization and growth improvement of *Avicennia marina*. *Aquat. Bot.*, 104: 101-105.
- Secilia, J. and D.J. Bagyaraj, 1992. Selection of efficient vesicular-arbuscular mycorrhizal fungi for wetland rice (*Oryza sativa* L.). *Biol. Fertil. Soils*, 13: 108-111.
- Secilia, J. and D.J. Bagyaraj, 1994. Selection of efficient vesicular-arbuscular mycorrhizal fungi for wetland rice: A preliminary screen. *Mycorrhiza*, 46: 265-268.
- Six, J., S.D. Frey, R.K. Thiet and K.M. Batten, 2006. Bacterial and fungal contributions to carbon sequestration in agro-ecosystems. *Soil Sci. Soc. Am. J.*, 70: 555-569.
- Shukla, N., R.P. Awasthi, R. Laxmi and J. Kumar, 2012. Biochemical and physiological responses of rice (*Oryza sativa* L.) as influenced by *Trichoderma harzianum* under drought stress. *Plant Physiol. Biochem.*, 54: 78-88.
- Smith, S.E., F.A. Smith and I. Jakobsen, 2003. Mycorrhizal fungi can dominate phosphate supply to plants irrespective of growth responses. *Plant Physiol.*, 133: 16-20.
- Solaiman, M.Z. and H. Hirata, 1997. Effect of arbuscular mycorrhizal fungi inoculation of rice seedlings at the nursery stage upon performance in the paddy field and greenhouse. *Plant Soil*, 191: 1-12.
- Sreenivasaprasad, S. and K. Manibhushanrao, 1990. Antagonistic potential of *Gliocladium virens* and *Trichoderma longibrachiatum* to phytopathogenic fungi. *Mycopathologia*, 109: 19-26.
- Tien, T.M., M.H. Gaskins and D.H. Hubbell, 1976. Plant growth substances produced by *Azospirillum brasilense* and their effect on the growth of pearl millet (*Pennisetum americanum* L.). *Appl. Environ. Microbiol.*, 37: 1016-1024.
- Tsavkelova, E.A., S. Yu, Klimova, T.A. Cherdyntseva and A.I. Netrusov, 2005. Microbial producers of plant growth stimulators and their practical use: A review. *Prikl. Biokhim. Mikrobiol.*, 42: 117-126.
- Vahed, H.S., P. Shahinrokhsar and F. Heydarnezhad, 2012. Performance of phosphate solubilizing bacteria for improving growth and yield of rice (*Oryza Sativa* L.) in the presence of phosphorus fertilizer. *Int. J. Agri. Crop. Sci.*, 4: 1228-1232.
- Vessey, J.K., 2003. Plant growth promoting rhizobacteria as biofertilizers. *Plant Soil*, 255: 571-586.
- Vyas, P. and A. Gulati, 2009. Organic acid production *in vitro* and plant growth promotion in maize under controlled environment by phosphate solubilizing *Fluorescent pseudomonas*. *BMC Microbiol.*, 9: 174.
- Wahyudi, A.T., R.I. Astuti and Giyanto, 2011. Screening of *Pseudomonas* sp. isolated from rhizosphere of soybean plant as plant growth promoter and biocontrol agent. *Am. J. Agr. Biol. Sci.*, 6: 134-141.
- Xie, C.H. and A. Yokota, 2006. *Zoogloea oryzae* sp. nov., a nitrogen-fixing bacterium isolated from rice paddy soil and reclassification of the strain ATCC 19623 as *Crabtreeella saccharophila* gen. nov., sp. nov. *Int. J. Syst. Evol. Microbiol.*, 56: 619-624.
- Xu, F., J. Hu, S. Wu and M.H. Wong, 2013. Grain yield and arsenic uptake of upland rice inoculated with arbuscular mycorrhizal fungi in as-spiked soils. *Environ. Sci. Pollut. Res. Int.*, DOI: 10.1007/s11356-012-1440-9.
- Yani, Y.G., R.Y. Rizk, F.K. Abd El-Fattah, A. Squartini, V. Corich, A. Giacomini, F. de Bruijn, J. Rademaker, J. Maya-Flores, P. Ostrom, M. Vega-Hernandez, R.I. Hollingsworth, E. Martinez-Molina, P. Mateos, E. Velazquez, J. Wopereis, E. Triplett, M. Umali-Garcia, J.A. Anarna, B.G. Rolfe, J.K. Ladha, J. Hill, R. Mujoo, P.K. Ng and F.B. Dazzo, 2000. The beneficial plant growth-promoting association of *Rhizobium leguminosarum* bv. *trifolii* with rice roots. *Aust. J. Plant Physiol.*, 28: 845-870.
- Yeasmin, T., P. Zaman, A. Rahman, N. Absar and N.S. Khanum, 2007. Arbuscular mycorrhizal fungus inoculum production in rice plants. *Afr. J. Agric. Res.*, 2: 463-467.
- Zhang, S.J., L. Wang, F. Ma, X. Zhang, Y.N. Xu and Z. Li, 2012. Effects of mycorrhiza alone with or without nitrogen and phosphate on rice dry matter production and distribution. *J. Harbin Inst. Technol.*, 44: 33-36.
- Zhang, X.H., Y.G. Zhu, B.D. Chen, A.J. Lin, S.E. Smith and F.A. Smith, 2005. Arbuscular mycorrhizal fungi contribute to resistance of upland rice to combined metal contamination of soil. *J. Plant Nutr.*, 28: 2065-2077.