

Research Article

Estrogen and Growth Hormone and their Roles in Reproductive Function

Hüseyin Baki ÇİFTÇİ

Department of Animal Science, School of Agriculture, Selçuk University, Konya 42075, Turkey

Abstract: The aim of this study was to review the effect of estrogen on growth hormone secretion and the roles of estrogen and growth hormone in reproductive function. Estrogen is the main hormone affecting growth, development, maturation and functioning of reproductive tract as well as the sexual differentiation and the behavior. Growth hormone is also important factor in sexual maturation and attainment of puberty. The impact of estrogen on growth hormone secretion has been reported in rodents and primates. However, the precise mechanism for the alterations in growth hormone secretion is not clearly known. Estrogen may possibly have a direct effect on growth hormone secretion via the binding to estrogen receptor- α due to its co-expression in growth hormone neurons in the medial preoptic area and arcuate nucleus. Estrogen may also have an indirect effect via the reducing insulin-like growth factor-1 feedback inhibition resulting with increased growth hormone secretion.

Keywords: Estrogen, growth, hormone, ovary, reproduction, testis

INTRODUCTION

Estrogen is an intra ovarian factor affecting the function of hypothalamus, pituitary, liver, skeleton and calcium homeostasis (Turner *et al.*, 1994). It is also a main reproductive hormone affecting growth, development, maturation and functioning of reproductive tract as well as the sexual differentiation and the behavior (Balthazart *et al.*, 2009). It was hypothesized that exogenous estrogens enhance GH secretion (Eden, 1979) and promote somatic growth. Therefore exogenous estrogens have been used to increase the secretory characteristics of GH in several species including sheep (Misztal *et al.*, 2007), cattle (Colak *et al.*, 2011) and human (Weissberger *et al.*, 1991). The main aim of this study was to review the effect of estrogen on growth hormone secretion and the roles of estrogen and growth hormone in reproductive function.

ESTROGEN

Estrogen is named for its importance in the estrous cycle. Animal body naturally produce three main forms of estrogen, which are estradiol 17β (E_2), Estrone (E_1) and Estriol (E_3). Estrone and estriol were firstly identified in the urine of pregnant women and this was followed by the identification of E_2 in the follicular fluid of sow by Edward Adelbert Doisy between 1929-1936 (Simoni *et al.*, 2002). Estradiol and estrone are formed, respectively, from testosterone and androstenedione by aromatase which catalyzes an aromatic hydroxylation of the A ring of C19 androgens. Estrone and estradiol are inter-converted by 17-

hydroxysteroid dehydrogenase; both can be converted to estriol by 16 α -hydroxylase, mainly in the liver (Moghrabi *et al.*, 1997).

In the ovary, E_2 is the most physiologically active type of estrogen produced by granulosa cells of pre-ovulatory follicles through the aromatization of thecal androgen by the granulosa cells of growing follicles. In the testis, some E_2 is produced by Sertoli cells through the aromatization of the androgen synthesized from the Leydig cells. Estrogen is also synthesized in extragonadal sites including the mesenchymal cells of adipose tissue and skin, osteoblasts and chondrocytes of bone, vascular endothelium and aortic smooth muscle cells as well as several sites in the brain, including the anterior hypothalamus and the medial basal hypothalamus (Bayard *et al.*, 1995; Sasano *et al.*, 1999).

Estrogen receptors: Estrogens act via two types of receptors (ER α and ER β), which are members of a large super family of proteins that function as ligand-activated transcription factors (Katzenellenbogen and Katzenellenbogen, 1996). Both receptors have direct differentiative influences on reproductive organs and have similar binding affinity to estradiol (Kuiper *et al.*, 1997; Drummond *et al.*, 1999). Although, there are significant amino acid differences in the regions of these receptors that would be expected to influence transcriptional activity (Hall and McDonnell, 1999). More recently, two Estrogen-Related Receptors, (ERR α /ERR1) and (ERR β /ERR2) have also been characterized.

The presence of oestrogen receptors have been shown within the hypothalamus, pituitary, ovary,

Table 1: Expression of estrogen receptors in reproductive organs of different animal species

	Hypothalamus	Pituitary	Ovary	Oviduct	Uterus	Cervix	Vagina
<i>F. primat</i>	+α/+β	+β	+α/+β	+α/+β	+α/+β	+α	+α/-β
<i>Ewe</i>	+α/+β	+α/+β	+α/+β	+α	+α/+β	+α	No data
<i>Cow</i>	+α/+β	+α/+β	+α/+β	+α/+β	+α/+β	+α	+α
<i>F. goat</i>	No data	+α/+β	+β	No data	+α	No data	No data
<i>Porcine</i>	+α/+β	+α/+β	+α/+β	+α	+α/+β	+α	No data
<i>F. rat</i>	+α/+β	+α/-β	+α/+β	+α/-β	+α/-β	+α/-β	+α/-β
<i>F. mouse</i>	+α/+β	+α/-β	+α/+β	+α/-β	+α/+β	+α/+β	+α/+β

Plus (+) indicates the presence while minus (-) indicates its absence; References are in the text




Fig. 1: Oestrogen receptor is composed of several functional regions

oviduct, uterus, cervix and vagina of several species including *human* (Brodowska *et al.*, 2007), *sheep* (Juengel *et al.*, 2006), *cow* (Sağsöz, 2011), *goat* (Cui *et al.*, 2009), *Porcine* (Knapczyk-Stwora *et al.*, 2011), *rat* (Okada *et al.*, 2003) and *mouse* (Hulaś-Stasiak and Gawron, 2007) (Table 1). Estrogen signaling is selectively stimulated or inhibited depending upon a balance between ERα and ERβ activities in target organs. ERs have five distinct regions (Skafar and Zhao, 2008). These distinct regions correspond to functional and structural units called domain (Fig. 1).

The N-terminal of the A and B domains are highly conserved only between chicken and human estrogen receptors, but this distinction is much less clear in the other steroid Receptors. Therefore, region A and B are combined into A/B region in most cases. This region is a modulatory region and it is the most variable both in size and sequence. The A/B domain consists of Activation Function 1 (AF1), which contributes to the transcriptional activity of ERs and is an essential domain for interaction with co-regulators. The C domain is DNA Binding Domain (DBD) and the most conserved region of the estrogen receptors. It is essential for sequence specific binding of ERs to DNA. The D domain is not well conserved among the different receptors and serves a hinge between the DBD and the Ligand Binding Domain (LBD) allowing rotation of the DBD. Domain E is the Ligand Binding Domain (LBD), which contains COOH-terminal AF-2 motif responsible for ligand-dependent transcriptional activation. The LBDs are folded into a three-layered, anti-parallel helical sandwich. A central core layer of three helices is packed between two additional layers to create a cavity. This cavity is completely partitioned from the external environment and is closed by helix 12 of the ligand binding domain, operating as a lid after hormone has entered binding pocket (Fig. 2). The relocation of helix 12 over the hormone binding site generates new surfaces that allow co-activator to bind ligand binding domain, thereby mediating the activity




Fig. 2: Ligand Binding Domain (LBD) of human oestrogen Receptor-α (ERα) complexed with the natural ligand 17β-oestradiol. A central core layer of three helices is packed between two additional layers to create a cavity. This cavity is completely partitioned from the external environment and is closed by helix 12 (Adopted from the Fig. 6 of Beato and Klug (2000)

of AF-2. The F region is essential for hormone binding in the ERα.

Effect of estradiol on reproductive function: Chromosomal sex is determined at the time of fertilization by the entry of an X or Y chromosome from the sperm pronucleus into the pronucleus of the oocyte (Marshall Graves, 2000). In accord to the chromosomal sex, gonads are formed. It was suggested that under the influence of the Y chromosome, the undifferentiated genital ridge develops into testis and in its absence, ovaries form by default. Therefore, the existence of two candidate genes (SRY and ZFY), former being confirmed as Testis Determining Factor (TDF), on the Y chromosome was predicted following the mutation analysis (Jost *et al.*, 1973; Marshall Graves, 2000).

Gonadal hormone secretion is under the control of chromosomal sex, which, in turn, controls the phenotype of non-gonadal tissue. The hormonal regulation of sexual differentiation of the mammalian reproductive system was established in the late 1940s by Jost. In his study, testes were removed from fetal male rabbits, inducing a female phenotype at birth. In contrast, transplantation of testis into female embryos

induced a male phenotype. The early fetus has the potential to be either male or female and possesses not only an undifferentiated gonadal ridge, but also 'precursors' for both Mullerian and Wolfian ducts. Once the testis are formed, they secrete Mullerian Inhibitory Substance (MIS), which induces regression of the Mullerian duct and they also produce testosterone, which stimulates development of the Wolfian duct. In the absence of testis and thus, MIS and testosterone, the Mullerian duct develops and the Wolfian duct regresses. Thus, it was established that male sexual development requires hormonal control and that the female reproductive system develops in the absence of these hormones (Jost *et al.*, 1973). Other studies have confirmed that estrogen has no effect on mullerian duct formation. Because, treatment of the pregnant mice with Diethylstilboestrol (DES) did not affect Müllerian duct formation in female embryos (Newbold and McLachlan, 1982). However, the presence of estrogen receptors in both male and female mice from gestational day 10 and later (Gorski and Hou, 1995) indicate that estrogen has role in reproductive tract development and functioning. According to a study, in mutant mice lacking responsiveness to estradiol by disrupting the estrogen receptor gene by gene targeting showed abnormal reproductive tract development. It was also noted that the males were infertile (Lubahn *et al.*, 1993). Experimental inhibition of the formation or action of estrogen in the female chicken and Japanese quail embryos can result in almost complete phenotypic sex-reversal, such as formation of testis-like ovaries, development of male secondary sex characteristics, lack of oviductal development and male-like growth of the cloacal gland in response to testosterone (Elbrecht and Smith, 1992).

The sex differences in the morphological and functional phenotype of the body and brain underlie gender identity, sexual orientation, sexual behavior and differences in certain non-reproductive behaviors. In most mammals, the principal hormone masculinizing the brain is testosterone. However, testosterone is the principal hormone causing brain masculinisation, but its metabolite, oestradiol, acting on estrogen receptors α and β (ER α and ER β) control separate aspects of differentiation. ER α is primarily involved in masculinization, while ER β mediates defeminization of sexual behaviors, but not masculinization (Kudwa *et al.*, 2006).

Estrogen also acts as an intra-gonadal factor and has negative and positive feedback influences on the hypothalamic-pituitary axis to regulate gonadotrophin secretion. It has been known for many years that estrogen has a direct influence on folliculogenesis. Oestradiol-17 β (E $_2$) and its analogues have both proliferative and differentiative effects on somatic cells of follicles (Findlay *et al.*, 2001). It stimulates the proliferation of granulosa cells in follicles and serves to facilitate the actions of Follicle Stimulating Hormone

(FSH) and Luteinizing Hormone (LH) (Richards, 1980). Thus, it permits follicle growth because, increase in follicle size is due directly to an increase in granulosa cell number and not due to the antrum formation (Goldenberg *et al.*, 1972). Estrogen is also responsible for facilitating the differentiation of granulosa cells including the induction of receptor systems for FSH, LH and prolactin and it can influence post-receptor mechanisms. There is a strong consensus that both ER α and ER β are expressed in granulosa cells of preantral and antral follicles (Drummond *et al.*, 1999). Estrogen Receptor- α Knockout (ERKO) female mice are acyclic, infertile and possess hyperemic ovaries devoid of corpora lutea (Couse and Korach, 1999). Folliculogenesis is arrested at the antral stage with large secondary follicles becoming cystic and hemorrhagic within 3 weeks of birth. In contrast, Estrogen Receptor-B Knockout (BERKO) females have small ovaries, some arrested follicular development and their fertility is compromised with reduced numbers of offspring per litter, consistent with the reduced number of corpora lutea observed (Krege *et al.*, 1998).

Estrogen is also synthesized in the male reproductive system and it is found in high concentrations in rete testis and seminal fluids. Both estrogen receptors (ER α and ER β) are found in various regions of the male reproductive tract. It was reported that estradiol (E $_2$) induces spermatogenesis in gonadotropin-deficient hypogonadal (*hpg*) mice (Allan *et al.*, 2010). It was concluded that E $_2$ -induced spermatogenesis in hypogonadal (*hpg*) mice involves an ER α -dependent neuroendocrine mechanism increasing blood FSH and Sertoli cell function (Allan *et al.*, 2010). The main breakthrough in this field was brought forth by estrogen receptor knockout mice. Phenotypically, these mice have significant alteration in testes histology, spermiogenesis and they suffer from infertility (Eddy *et al.*, 1996).

GROWTH HORMONE STRUCTURE AND SIGNALING

Growth Hormone (GH) is an anti-parallel four-helix bundle protein (Chantalat *et al.*, 1995) secreted in a pulsatile manner by somatotrophs in the anterior pituitary gland (Edmondson *et al.*, 2003). Its secretion is controlled by two neuropeptides namely Growth Hormone-Releasing Hormone (GHRH) and the Somatostatin (SS). Somatostatin (SS) inhibits GH release without affecting GH synthesis. Several lines of evidence suggest that GHRH initiates GH pulses and somatostatin modulates the amplitude of GH pulses. Blocking the action of GHRH, either by passive immunization in rats or with a GHRH antagonist in rats or humans, abolishes pulsatile GH release. Growth hormone also exerts a negative feedback effect on its own secretion. Daily subcutaneous administration of GH for 2-5 days decrease the endogenous GH response




Fig. 3: The crystal structure of the human GH and extracellular domain of its receptor. GH has two distinct binding faces to GHR on either side of the molecule, known as site 1 and 2. The extracellular domain of GHR comprises two beta sandwiches named domain GHR1 and GHR2

to GHRH (Ross *et al.*, 1987). This effect may be mediated through the secretion of IGF from the liver.

The anterior pituitary gland synthesizes different GH isoforms. In human, the genetic locus encoding GH resides in long arm of chromosome 17 (17q24.2.) Two genes, in the cluster, encode two distinct GH variants and are named GH1 (or GH-N) and GH2 (or GH-V). GH1 is the principal and most abundant GH form in the pituitary. The GH2 gene product, GH2 (or GH-V) is similar in structure to the GH1. Its sequence differs from that of GH1 at 13 amino acid positions. Two other isoforms are Chorionic Somatomammotropin (CS), also known as Placental Lactogen (PL) named as CS1 (or CSA) and CS2 (or CSB) expressed in the placenta.

Growth hormone has two distinct binding faces to its receptor (GHR) on either side of the molecule, known as site 1 and 2, which are buried by essentially identical receptor domains upon binding (De Vos *et al.*, 1992). The first binding site (site 1) has a concave shape and is formed by residues that are exposed on helix 4 of the helix bundle, together with residues on the connecting loop between helix 1 and 2 to produce an extensive binding crevice (De Vos *et al.*, 1992). The second binding site (site 2) is made-up of the exposed sites of helices 1 and 3. In contrast to the concave character of site 1, site 2 is relatively flat. The hormone interacts through site-1 with the first molecule of GHR and promotes the dimerization of the receptor by interaction of the lower-affinity binding site 2 (Fig. 3).

Upon binding, GH causes dimerization of GHR, activation of the GHR-associated JAK2 tyrosine kinase and tyrosyl phosphorylation of both JAK2 and GHR. These events recruit and/or activate a variety of signaling molecules, including MAP kinases, insulin receptor substrates, phosphatidylinositol 3' phosphate kinase, diacylglycerol, protein kinase C, intracellular calcium and STAT transcription factors. These signaling molecules contribute to the GH-induced changes in enzymatic activity, transport function and gene expression that ultimately culminate in changes in growth and metabolism.

Effect of growth hormone on reproductive function:

Growth Hormone (GH) is important factor in sexual maturation and attainment of puberty. External administration of GH has been shown to accelerate sexual maturation in monkeys (Wilson *et al.*, 1989) and GH-deficient children (Stanhope *et al.*, 1992). GH may accelerate puberty by activating the Luteinizing Hormone (LH) -releasing hormone pulse generator (Bartke *et al.*, 1999) and/or by potentiating androgen action (Ilondo *et al.*, 1982). However, GH administration has not been shown to accelerate pubertal development in pigs (Andres *et al.*, 1991). In human, GH appears to increase the rate of human sexual maturation only when a pubertal pattern of pituitary gonadotrophin secretion is established (Sharara and Giudice, 1997). This shows that effect of external GH administration on sexual maturation depends on the reproductive state at the time of GH treatment. In young female rat, the injection of bovine GH (bGH) to the median eminence delays puberty (Advis *et al.*, 1981), it is possible that GH exerts inhibitory effects on the hypothalamo-pituitary-gonadal axis at central sites, contrary to its stimulatory actions on pituitary gonadal function.

Growth hormone gene is also expressed within the gonads. By using cDNA primers for pituitary GH in Reverse Transcription-Polymerase Chain Reaction (RT-PCR), GH mRNA was shown within testis of fetal mice (Nguyen *et al.*, 1996), adult cockerels (Harvey *et al.*, 2004; Luna *et al.*, 2004) and human (Berger *et al.*, 1999). By using in situ hybridization and immunohistochemistry, GH mRNA and protein expression were shown in the ovary of hen (Ahumada-Solórzano *et al.*, 2012). In the chicken testis, the GH-immunoreactivity was not present in spermatogonia, but it was mainly in the primary and secondary spermatocytes and spermatids and in the luminal compartments of Sertoli cells, but was also in surrounding myocytes and interstitial cells. The localization of GH in spermatocytes and spermatids suggests unsuspected roles in gamete development.

Within the ovary, GH play important role in early, Follicle-Stimulating Hormone (FSH) -independent follicular development, since GH-binding activity peaks

during early folliculogenesis in porcine follicles (Quesnel, 1999) and fish ovarian homogenates (Gomez *et al.*, 1998). *In vivo* and *in vitro* studies suggest that GH stimulates growth and prevents atresia in small follicles. Growth hormone is important factor in follicle recruitment and initiation of oocyte growth. It acts together with gonadotrophins to stimulate later stages of folliculogenesis and luteinization, since both GH and gonadotrophins are required to prevent atresia of larger follicles (>2 mm) following hypophysectomy in sheep (Eckery *et al.*, 1997). GH administration *in vivo* increases the number of large follicles in pigs (Lucy *et al.*, 1995), GH-deficient dwarf rats (Ozawa *et al.*, 1996) and the number of corpora lutea in cattle (Lucy *et al.*, 1992). GH may play a role in follicle selection, since GH-binding sites in sow granulosa cells are lost in atretic follicles (Quesnel, 1999) and the development of the dominant follicle is impaired in GHR deficient cattle (Chase *et al.*, 1998).

GH may facilitate ovulation by increasing sensitivity to gonadotrophins and by reducing the incidence of apoptosis in preovulatory ovarian follicles. The increased number of corpora lutea and reduced numbers of atretic follicles in the ovaries of mice transgenically expressing GH supports this view (Danilovich *et al.*, 2000). The over expression of GH in these mice has thus been correlated with an increase in the number of ova shed during each ovulation (Danilovich *et al.*, 2000).

Effect of estrogen on growth hormone secretion:

Estrogen effect growth and reproduction at numerous physiological levels at target sites, including direct actions at the hypothalamic and pituitary levels to modulate Growth Hormone (GH) production and secretion. In ovary-intact mice GH mRNA in the arcuate nucleus and Medial Preoptic Area (MPOA) had elevated, while ovariectomy decreased GH mRNA in both regions (Addison and Rissman, 2012). When gonadectomized adults of both sexes were treated with estradiol, GH mRNA increased in females but had no effect in castrated males. It was also found that estrogen receptor- α is co-expressed in GH neurons in the MPOA and arcuate nucleus (Addison and Rissman, 2012). In man, estrogen produced locally from aromatization plays a major role in the regulation of GH secretion (Birzniece *et al.*, 2010). It was proposed that exogenous estrogens enhance GH secretion (Eden, 1979). In women, the effect of exogenous estrogen on the regulation of GH secretion is route-dependent. Oral administration of estrogen enhances GH secretion; however, this does not happen when estrogen is replaced by a physiological non-oral route (Weissberger *et al.*, 1991). There is no consensus on a mechanism (especially in ruminants) for alterations in the GH axis by estrogenic substances (Carroll *et al.*, 2007). Since not all estrogens have the same effect on pulsatile GH,

the dose of estradiol may influence the type of effects of estradiol on GH pulses.

Estrogens may also indirectly stimulate GH secretion by reducing IGF-I feedback inhibition through IGF-I modulation. IGF-I mediates a negative feedback control of GH secretion by acting directly on hypothalamic GHRH and Somatostatin (SS) neurons that the route of administration is a major determinant of the effect of estrogen on the GH/IGF-I axis (Leung *et al.*, 2004).

CONCLUSION

Both estrogen and growth hormone have major impacts on steroidogenesis, gametogenesis, gonadal differentiation as well as the attainment of sexual maturity. Estrogen modulates growth hormone secretion via the different pathways Such as an effect on hypothalamic neurons and/or directly via effect on pituitary somatotrops also and/or indirectly via the reducing IGF-I feedback inhibition. The effect of external estrogen administration depends on dose, route of administration, species, gender and age. The presence of multi GH isoforms presents difficulties for an exact measurement of the hormone in body fluids and for the clear understanding of its physiology.

REFERENCES

- Addison, M.L. and E.F. Rissman, 2012. Sexual dimorphism of growth hormone in the hypothalamus: Regulation by estradiol. *Endocrinology*, 153(4): 1898-1907.
- Advis, J.P., S.S. White and S.R. Ojeda, 1981. Activation of growth hormone short loop negative feedback delays puberty in the female rat. *Endocrinology*, 108(4): 1343-1352.
- Ahumada-Solórzano, S.M., M.E. Carranza, E. Pedernera, A.J. Rodríguez-Méndez, M. Luna and C. Arámburo, 2012. Local expression and distribution of growth hormone and growth hormone receptor in the chicken ovary: Effects of GH on steroid genesis in cultured follicular granulosa cells. *Gen. Comp. Endocrinology*, 175(2): 297-310.
- Allan, C.M., J.F. Couse, U. Simanainen, J. Spaliviero, M. Jimenez, K. Rodriguez, K.S. Korach and D.J. Handelsman, 2010. Estradiol induction of spermatogenesis is mediated via an estrogen receptor- α mechanism involving neuroendocrine activation of follicle-stimulating hormone secretion. *Endocrinology*, 151(6): 2800-2810.
- Andres, C.J., M.L. Green, J.A. Clapper, T.R. Cline and M.A. Diekman, 1991. Influence of daily injections of porcine somatotropin on growth, puberty and reproduction in gilts. *J. Anim. Sci.*, 69(9): 3754-3761.

- Balthazart, J., C.A. Cornil, T.D. Charlier, M. Taziaux and G.F. Ball, 2009. Estradiol, a key endocrine signal in the sexual differentiation and activation of reproductive behavior in quail. *J. Exp. Zool. Ecol. Genet. Physiol.*, 311(5): 323-345.
- Bartke, A., V. Chandrashekar, D. Turyn, R.W. Steger, L. Debeljuk, T.A. Winters, J.A. Mattison, N. Danilovich, W. Croson, D.R. Wernsing and J. Kopchick, 1999. Effects of growth hormone overexpression and growth hormone resistance on neuroendocrine and reproductive functions in transgenic and knock-out mice. *Proc. Soc. Exp. Biol. Med.*, 222(2): 113-123.
- Bayard, F., S. Clamens, G. Delsol, N. Blaes, A. Maret and J.C. Faye, 1995. Oestrogen biosynthesis, oestrogen metabolism and functional oestrogen receptors in bovine aortic endothelial cells. *Ciba Found. Symp.*, 191: 122-132.
- Beato, M. and J. Klug, 2000. Steroid hormone receptors: An update. *Hum. Reprod. Update*, 6(3): 225-236.
- Berger, P., G. Untergasser, M. Hermann, A. Hittmair, S. Madersbacher and S. Dirnhofer, 1999. The testis-specific expression pattern of the Growth Hormone/Placental Lactogen (GH/PL) gene cluster changes with malignancy. *Hum. Pathol.*, 30(109): 1201-1206.
- Birzniece, V., A. Sata, S. Sutanto and K.K. Ho, 2010. Paracrine regulation of growth hormone secretion by estrogen in women. *J. Clin. Endocrinol. Metab.*, 95(8): 3771-3776.
- Brodowska, A., M. Laszczynska, A. Starczewski, B. Karakiewicz and J. Brodowski, 2007. The localization of estrogen receptor alpha and its function in the ovaries of postmenopausal women. *Folia Histochem. Cytobiol.*, 45(4): 325-330.
- Carroll, J.A., M.A. Walker, S.M. Hartsfield, N.H. McArthur and T.H. Welsh, 2007. Visual documentation of ovine pituitary gland development with magnetic resonance imaging following zeranol treatment. *Lab. Anim.*, 41(19): 120-127.
- Chantalat, L., N.D. Jones, F. Korber, J. Navaza and A. Pavlovsky, 1995. The crystal structure of wild-type growth-hormone at 2.5 Å resolution. *Protein Pept. Lett.*, 2: 333-340.
- Chase, C.C., C.J. Kirby, A.C. Hammond, T.A. Olson and M.C. Lucy, 1998. Patterns of ovarian growth and development in cattle with a growth hormone receptor deficiency. *J. Anim. Sci.*, 76(1): 212-219.
- Colak, M., T. Shimizu, N. Matsunaga, C. Murayama, S. Nagashima, M. Kataoka, C. Kawashima, M. Matsui, H.A. van Dorland, R.M. Bruckmaier and A. Miyamoto, 2011. Oestradiol enhances plasma growth hormone and insulin-like growth factor-I concentrations and increased the expression of their receptors mRNAs in the liver of ovariectomized cows. *Reprod. Domest. Anim.*, 46(5): 854-861.
- Couse, J.F. and K.S. Korach, 1999. Reproductive phenotypes in the estrogen receptor-alpha knockout mouse. *Ann. Endocrinol. (Paris)*, 60(2): 143-148.
- Cui, H.X., S.M. Zhao, M.L. Cheng, L. Guo, R.Q. Ye, W.Q. Liu and S.Z. Gao, 2009. Cloning and expression levels of genes relating to the ovulation rate of the Yunling black goat. *Biol. Reprod.*, 80(2): 219-226.
- Danilovich, N., A. Bartke and T.A. Winters, 2000. Ovarian follicle apoptosis in bovine growth hormone transgenic mice. *Biol. Reprod.*, 62(1): 103-107.
- De Vos, A.M., M. Ultsch and A.A. Kossiakoff, 1992. Human growth hormone and extracellular domain of its receptor: Crystal structure of the complex. *Science*, 255(5042): 306-312.
- Drummond, A.E., A.J. Baillie and J.K. Findlay, 1999. Ovarian estrogen receptor alpha and beta mRNA expression: Impact of development and estrogen. *Mol. Cell. Endocrinol.*, 149(1-2): 153-161.
- Eckery, D.C., C.L. Moeller, T.M. Nett and H.R. Sawyer, 1997. Localization and quantification of binding sites for follicle-stimulating hormone, luteinizing hormone, growth hormone and insulin-like growth factor I in sheep ovarian follicles. *Biol. Reprod.*, 57(39): 507-513.
- Eddy, E.M., T.F. Washburn, D.O. Bunch, E.H. Goulding, B.C. Gladen, D.B. Lubahn and K.S. Korach, 1996. Targeted disruption of the estrogen receptor gene in male mice causes alteration of spermatogenesis and infertility. *Endocrinology*, 137(11): 4796-4805.
- Eden, S., 1979. Age and sex-related differences in episodic growth hormone secretion in the rat. *Endocrinology*, 105(2): 555-560.
- Edmondson, S.R., S.P. Thumiger, G.A. Werther and C.J. Wraight, 2003. Epidermal homeostasis: The role of growth hormone and insulin-like growth factor systems. *Endocr. Rev.*, 24(6): 737-764.
- Elbrecht, A. and R.G. Smith, 1992. Aromatase enzyme activity and sex determination in chickens. *Science*, 255(5043): 467-470.
- Goldenberg, R.L., J.L. Vaitukaitis and G.T. Ross, 1972. Estrogen and follicle-stimulating hormone interactions on follicle growth in rats. *Endocrinology*, 90(6): 1492-1498.
- Gomez, J.M., M. Loir and F. Le Gac, 1998. Growth hormone receptors in testis and liver during the spermatogenic cycle in rainbow trout (*Oncorhynchus mykiss*). *Biol. Reprod.*, 58(2): 483-491.
- Gorski, J. and Q. Hou, 1995. Embryonic estrogen receptors: Do they have a physiological function? *Environ. Health Perspect.*, 103(Suppl 7): 69-72.
- Hall, J.M. and D.P. McDonnell, 1999. The estrogen receptor β -isoform (ER β) of the human estrogen receptor modulates ER α transcriptional activity and is a key regulator of the cellular response to estrogens and antiestrogens. *Endocrinology*, 140(12): 5566-5578.

- Harvey, S., M.L. Baudet, A. Murphy, M. Luna, K.L. Hull and C. Aramburo, 2004. Testicular Growth Hormone (GH): GH expression in spermatogonia and primary spermatocytes. *Gen. Comp. Endocrinol.*, 139(2): 158-167.
- Hułas-Stasiak, M. and A. Gawron, 2007. Immunohistochemical localization of estrogen receptors ERalpha and ERbeta in the spiny mouse (*Acomys cahirinus*) ovary during postnatal development. *J. Mol. Histol.*, 38(1): 25-32.
- Ilonzo, M.M., M. Vanderschueren-Lodeweyckx, R. Vlietnick, M. Pizarro, P. Malvaux, E. Eggermont and R. Eeckels, 1982. Plasma androgens in children and adolescents Part II: A longitudinal study in patients with hypopituitarism. *Horm. Res.*, 16(2): 78-95.
- Jost, A., B. Vigier, J. Prepin and J.P. Perchellet, 1973. Studies on sex differentiation in mammals. *Recent. Prog. Horm. Res.*, 29: 1-41.
- Juengel, J.L., D.A. Heath, L.D. Quirke and K.P. McNatty, 2006. Oestrogen receptor alpha and beta androgen receptor and progesterone receptor mRNA and protein localisation within the developing ovary and in small growing follicles of sheep. *Reproduction*, 131(1): 81-92.
- Katzenellenbogen, J.A. and B.S. Katzenellenbogen, 1996. Nuclear hormone receptors: Ligand activated regulators of transcription and diverse cell responses. *Chem. Biol.*, 3(7): 529-536.
- Knapczyk-Stwora, K., M. Durlej, M. Duda, K. Czernichowska-Ferreira, A. Tabecka-Lonczynska and M. Slomczynska, 2011. Expression of oestrogen receptor α and oestrogen receptor β in the uterus of the pregnant swine. *Reprod. Domest. Anim.*, 46(1): 1-7.
- Krege, J.H., J.B. Hodgins, J.F. Couse, E. Enmark, M. Warner, J.F. Mahler, M. Sar, K.S. Korach, J.A. Gustafsson and O. Smithies, 1998. Generation and reproductive phenotypes of mice lacking estrogen receptor beta. *PNAS*, 95(26): 15677-15682.
- Kudwa, A.E., V. Michopoulos, J.D. Gatewood and E.F. Rissman, 2006. Roles of estrogen receptors α and β in differentiation of mouse sexual behavior. *Neuroscience*, 138(3): 921-928.
- Kuiper, G.G., B. Carlsson, K. Grandien, E. Enmark, J. Haggblad, S. Nilsson and J.A. Gustafsson, 1997. Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptors alpha and beta. *Endocrinology*, 138(3): 863-870.
- Leung, K.C., G. Johannsson, G.M. Leong and K.K. Ho, 2004. Estrogen regulation of growth hormone action. *Endocr. Rev.*, 25(5): 693-721.
- Lubahn, D.B., J.S. Moyer, T.S. Golding, J.F. Couse, K.S. Korach and O. Smithies, 1993. Alteration of reproductive function but not prenatal sexual development after insertional disruption of the mouse estrogen receptor gene. *Proc. Natl. Acad. Sci. U.S.A.*, 90(23): 11162-11166.
- Lucy, M.C., W.W. Thatcher, R.J. Collier, F.A. Simmen, Y. Ko, J.D. Savio and L. Badinga, 1995. Effects of somatotropin on the conceptus, uterus and ovary during maternal recognition of pregnancy in cattle. *Domest. Anim. Endocrinol.*, 12(1): 73-82.
- Lucy, M.C., W.W. Thatcher, J.D. Savio, G. Danet-Desnoyers, M.T. Moser, L. Badinga, F.A. Simmen and R.J. Collier, 1992. Effect of bovine somatotropin on ovarian follicles, corpora lutea and embryos during early pregnancy in cattle. *J. Anim. Sci.*, 70: Abstract no 271.
- Luna, M., L. Huerta, L. Berumen, H. Martínez-Coria, S. Harvey and C. Arámburo, 2004. Growth hormone in the male reproductive tract of the chicken: Heterogeneity and changes during ontogeny and maturation. *Gen. Comp. Endocrinol.*, 137(1): 37-49.
- Marshall Graves, J.A., 2000. Human Y chromosome, sex determination and spermatogenesis: A feminist view. *Biol. Reprod.*, 63(3): 667-676.
- Misztal, T., M. Wańkowska, K. Górski and K. Romanowicz, 2007. Central estrogen-like effect of genistein on growth hormone secretion in the ewe. *Acta. Neurobiol. Exp. (Wars)*, 67(4): 411-419.
- Moghrabi, N., J.R. Head and S. Andersson, 1997. Cell type-specific expression of 17 β -hydroxysteroid dehydrogenase type 2 in human placenta and fetal liver. *J. Clin. Endocrinol. Metab.*, 82(11): 3872-3878.
- Newbold, R.R. and J.A. McLachlan, 1982. Vaginal adenosis and adenocarcinoma in mice exposed prenatally or neonatally to diethylstilbestrol. *Cancer Res.*, 42(5): 2003-2011.
- Nguyen, A.P., A. Chandorkar and C. Gupta, 1996. The role of growth hormone in fetal mouse reproductive function. *Endocrinology*, 137(9): 3659-3666.
- Okada, A., Y. Ohta, S. Inoue, H. Hiroi, M. Muramatsu and T. Iguchi, 2003. Expression of estrogen, progesterone and androgen receptors in the oviduct of developing, cycling and pre-implantation rats. *J. Mol. Endocrinol.*, 30: 301-315.
- Ozawa, K., H. Mizunuma, H. Ozawa and Y. Ibuki, 1996. Recombinant human growth hormone acts on intermediate-sized follicles and rescues growing follicles from atresia. *Endocr. J.*, 43(1): 87-92.
- Quesnel, H., 1999. Localization of binding sites for IGF-I, insulin and GH in the sow ovary. *J. Endocrinol.*, 163(2): 363-372.
- Richards, J.S., 1980. Maturation of ovarian follicles: Actions and interactions of pituitary and ovarian hormones on follicular cell differentiation. *Physiol. Rev.*, 60(1): 51-89.

- Ross, R.J., F. Borges, A. Grossman, R. Smith, L. Ngahfoong, L.H. Rees, M.O. Savage and G.M. Besser, 1987. Growth hormone pretreatment in man blocks the response to growth hormone-releasing hormone; evidence for a direct effect of growth hormone. *Clin. Endocrinol. (Oxf)*, 26(1): 117-123.
- Sağsöz, H., M.E. Akbalik, B.G. Saruhan and M.A. Ketani, 2011. Localization of estrogen receptor α and progesterone receptor B in bovine cervix and vagina during the follicular and luteal phases of the sexual cycle. *Biotech. Histochem.*, 86(4): 262-271.
- Sasano, H., H. Murakami, S. Shizawa, S. Satomi, H. Nagura and N. Harada, 1999. Aromatase and sex steroid receptors in human vena cava. *Endocr. J.*, 46(2): 233-242.
- Sharara, F.I. and L.C. Giudice, 1997. Role of growth hormone in ovarian physiology and onset of puberty. *J. Soc. Gynecol. Investig.*, 4(1): 2-7.
- Simoni, R.D., R.L. Hill and M. Vaughan, 2002. The discovery of estrone, estriol and estradiol and the biochemical study of reproduction: The work of Edward Adelbert Doisy. *J. Biol. Chem.*, 277(28): 17.
- Skafar, D.F. and C. Zhao, 2008. The multifunctional estrogen receptor-alpha F domain. *Endocrine*, 33(1): 1-8.
- Stanhope, R., A. Albanese, P. Hindmarsh and C.G. Brook, 1992. The effects of growth hormone therapy on spontaneous sexual development. *Horm. Res.*, 38(Suppl 1): 9-13.
- Turner, R.T., B.L. Riggs and T.C. Spelsberg, 1994. Skeletal effects of estrogen. *Endocr. Rev.*, 15(3): 275-300.
- Weissberger, A.J., K.K. Ho and L. Lazarus, 1991. Contrasting effects of oral and transdermal routes of estrogen replacement therapy on 24-hour Growth Hormone (GH) secretion, insulin-like growth factor I and GH-binding protein in postmenopausal women. *J. Clin. Endocrinol. Metab.*, 72(2): 374-381.
- Wilson, M.E., T.P. Gordon, C.G. Rudman and J.M. Tanner, 1989. Effects of growth hormone on the tempo of sexual maturation in female rhesus monkeys. *J. Clin. Endocrinol. Metab.*, 68(1): 29-38.